

## Danshen: a phytochemical and pharmacological overview

MEI Xiao-Dan<sup>1△</sup>, CAO Yan-Feng<sup>1△</sup>, CHE Yan-Yun<sup>2</sup>, LI Jing<sup>3</sup>, SHANG Zhan-Peng<sup>1</sup>,  
ZHAO Wen-Jing<sup>1</sup>, QIAO Yan-Jiang<sup>1\*</sup>, ZHANG Jia-Yu<sup>4\*</sup>

<sup>1</sup> School of Chinese Pharmacy, Beijing University of Chinese Medicine, Beijing 102488, China;

<sup>2</sup> College of Pharmaceutical Science, Yunnan University of Traditional Chinese Medicine, Kunming 650500, China;

<sup>3</sup> College of Basic Medicine, Jinzhou Medical University, Jinzhou 121001, China;

<sup>4</sup> Beijing Research Institute of Chinese Medicine, Beijing University of Chinese Medicine, Beijing 100029, China

Available online 20 Jan., 2019

**[ABSTRACT]** Danshen, the dried root or rhizome of *Salvia miltiorrhiza* Bge., is a traditional and folk medicine in Asian countries, especially in China and Japan. In this review, we summarized the recent researches of Danshen in traditional uses and preparations, chemical constituents, pharmacological activities and side effects. A total of 201 compounds from Danshen have been reported, including lipophilic diterpenoids, water-soluble phenolic acids, and other constituents, which have showed various pharmacological activities, such as anti-inflammation, anti-oxidation, anti-tumor, anti-atherogenesis, and anti-diabetes. This article intends to provide novel insight information for further development of Danshen, which could be of great value to its improvement of utilization.

**[KEY WORDS]** Danshen; Traditional uses; Chemical constituents; Quality control; Pharmacological activities

**[CLC Number]** R965 **[Document code]** A **[Article ID]** 2095-6975(2019)01-0059-22

### Introduction

Medicinal herbal products have been used for healthcare in Asia for over a millennium, and the usage has continued in modern society. Many Chinese medicinal herbal derivatives have been introduced into global market in the past decades. Among the well-known Chinese herbal medicines, Danshen, the dried root or rhizome of *Salvia miltiorrhiza* Bge., plays an important role in maintaining the well-being of Chinese population. It is commonly utilized for improving body function, such as promoting blood circulation and restoring/enhancing immunity. According to the phytochemical studies, active compounds of the processed herb are divided into two groups: diterpenoids and phenolic acids. The chemical constituents possess several therapeutic effects, such as improving micro-circulation, anti-atherosclerosis, anti-inflammation, anti-tumor and alleviating diabetes.

Although several literatures on the chemical constituents and biological activities of Danshen have been published, these publications are not comprehensive. More constituents of Danshen have been isolated and identified, and more information on therapeutic uses has been acquired since the publication of literatures on similar topic. Hence, this review is constructed to provide an up-to-date information on Danshen, including its chemical constituents, pharmacology, traditional uses and preparations, modern pharmaceutical products and side effects in clinic. Its quality control is also discussed. The goal is to provide a valuable and comprehensive reference for further development and utilization of Danshen.

### Traditional Uses

Danshen, which is characterized by a series of functions, has been used in traditional Chinese medicine for approximately two millennia. The functions of Danshen summarized from clinical observations include promoting blood circulation to remove blood stasis, nourishing blood to tranquillize the mind, cooling blood to disperse carbuncles. The processed herb is used to treat various pains caused by blood stasis, lumps in the chest and abdomen, pyogenic infection and carbuncle of the skin, palpitation and insomnia. Danshen was originally recorded in Shennong Bencao Jing (the first Chinese Materia Medica in China during the Eastern Han Dynasty, 25 AD–220 AD). In

**[Received on]** 17-Oct.-2018

**[Research funding]** This work was supported by the National Natural Science Foundation of China (No. 81430094) and Beijing Nova Program (No. Z171100001117029).

**[\*Corresponding author]** E-mails: yjqiao@263.Net (QIAO Yan-Jiang); zhangjiayu0615@163.com (ZHANG Jia-Yu)

<sup>△</sup>The first two authors contributed equally to this work.

These authors have no conflict of interest to declare.

additional to aforementioned functions, it was used for external applications as well. Mingyi Bielu (Han Dynasty, 219 AD) has also stated that the herb could treat stiffness along the spinal column and feet numbness. Wupu Bencao (Wei and Jin Dynasties, 420 AD–589 AD) has recorded this herb for the treatment of pain in the chest and abdomen. Moreover, it has also described the medical plant in detail, including habitats, shape, eco-environment, harvest, storage and functions. Furthermore, a very clear and detailed record of the plant morphology and medicinal characters was kept in Shu Bencao (Five Dynasties and Ten States periods, 935 AD–960 AD). In addition, its functions, meridian attribution and many other aspects were documented in a comprehensive and concrete record in Bencao Gangmu (Ming Dynasty, 1578 AD). At present, the functions of Danshen, activating blood circulation to remove blood stasis, promoting menstrual discharge to relieve menalgia, clearing heart heat to relieve restlessness and cooling blood to disperse carbuncles, have been stated in official record, the *Chinese Pharmacopoeia* (2015).

In the course of Sino-foreign cultural exchanges, the medicinal uses of Danshen spread abroad, thereafter has been

duly recorded locally. YI Xin-Fang (Japan, 982 AD) and Dongyi Baojian (Korea, 1611AD) included accounts on Danshen. Nowadays, products containing Danshen are sold commercially for promoting circulation and alleviating blood stasis in Japan [1]. Similar products are also available in natural health retailers in American and European countries [1].

As it is well-known, Danshen is one of the oldest and most frequently used herbal medicines in traditional preparations, which are mainly formulated as a decoction. For instance, Danshen Decoction could be used for treating chill and fever of children, Haitongpi Decoction was used to treat wind-dampness, while Qingying Decoction was indicated for the treatment of acute infectious febrile diseases. Nowadays, the preparations of Danshen have been widely used in clinic for various diseases in China. For example, Fufang Danshen tablet is used to treat chest pain caused by angina pectoris, Guanxin Danshen capsule and Fufang Danshen Dripping pill have therapeutic effects on treating chest impediment syndrome due to qi-stagnation and blood-stasis. The main traditional and modern uses of Danshen in China are listed in Table 1.

**Table 1 The main traditional and modern uses of Danshen in China**

Preparation Name	Compositions	Clinic uses	References
Danshen Mogao	<i>Salvia miltiorrhiza</i> Bge., <i>Omphalia lapidescens</i> Schroet.	Treating epilepsy induced by terror and fever	<i>Qian Jin Fang</i> (Tang Dynasty, AD 652)
Danshen Pill	<i>Salvia miltiorrhiza</i> Bge., <i>Eucommia ulmoides</i> Oliv., <i>Achyranthes bidentata</i> Bl., <i>Dipsacus asper</i> Wall. ex Henry, <i>Cinnamomum cassia</i> Presl, <i>Zingiber officinale</i> Rosc.	Treating pain in the loins	<i>Sheng Ji Zong Lu</i> (Song Dynasty, AD 1117)
Danshen Decoction	<i>Salvia miltiorrhiza</i> Bge., <i>Sophora flavescens</i> Ait., <i>Cynanchum atratum</i> Bge., <i>Cinnamomum cassia</i> Presl, <i>Acorus tatarinowii</i> Schott, <i>Omphalia lapidescens</i> Schroet., <i>Cnidium monnieri</i> (L.) Cuss.	Treating chill and fever of children	<i>Pu Ji Fang</i> (Ming Dynasty, AD 1390)
Danshen San	<i>Salvia miltiorrhiza</i> Bge.	Treating menstrual disorders and threatened abortion	<i>Fu Ren Liang Fang</i> (the Southern Song Dynasty, AD 1237)
Danshen Yinzi	<i>Salvia miltiorrhiza</i> Bge., <i>Angelica sinensis</i> (Oliv.) Diels, <i>Atractylodes macrocephala</i> Koidz., <i>Asparagus cochinchinensis</i> (Lour.) Merr., <i>Ophiopogon japonicus</i> (L.f) Ker-Gawl., <i>Fritillaria cirrhosa</i> D.Don, <i>Citrus reticulata</i> Blanco, <i>Anemarrhena asphodeloides</i> Bge., <i>Glycyrrhiza uralensis</i> Fisch., <i>Acorus tatarinowii</i> Schott, <i>Coptis chinensis</i> Franch., <i>Schisandra chinensis</i> (Turcz.) Baill.	Treating amnesia	<i>Gu Jin Yi Tong</i> (Ming Dynasty, AD 1556)
Haitongpi Decoction	<i>Erythrina variegata</i> L. var. <i>orientalis</i> (L.) Merr., <i>Salvia miltiorrhiza</i> Bge., <i>Stephania tetrandra</i> S. Moore, <i>Glycyrrhiza uralensis</i> Fisch., <i>Ephedra sinica</i> Stapf, <i>Asparagus cochinchinensis</i> (Lour.) Merr., <i>Aconitum carmichaelii</i> Debx., <i>Cinnamomum cassia</i> Presl, <i>Zingiber officinale</i> Rose.	Treating wind-dampness	<i>Sheng Ji Zong Lu</i> (Song Dynasty, AD 1117)
Tianwang Buxin Dan	<i>Rehmannia glutinosa</i> Libosch., <i>Panax ginseng</i> C. A. Mey., <i>Salvia miltiorrhiza</i> Bge., <i>Scrophularia ningpoensis</i> Hemsl., <i>Poria cocos</i> (Schw.) Wolf, <i>Polygala tenuifolia</i> Willd., <i>Platycodon grandiflorum</i> (Jacq.) A.DC., <i>Schisandra chinensis</i> (Turcz.) Baill., <i>Angelica sinensis</i> (Oliv.) Diels, <i>Asparagus cochinchinensis</i> (Lour.) Merr., <i>Ophiopogon japonicus</i> (L.f) Ker-Gawl., <i>Platycladus orientalis</i> (L.) Franco, <i>Ziziphus jujuba</i> Mill. var. <i>spinosa</i> (Bunge) Hu ex H. F. Chou	Tonifying the blood and nourishing the heart to cause tranquilization	<i>She Sheng Mi Pou</i> (Ming Dynasty, AD 1638)
Tiaojing Pill	<i>Salvia miltiorrhiza</i> Bge.	Treating irregular menstruation, dysmenorrhea, metrorrhagia and leukorrhagia due to stagnation of qi and blood	<i>Ji Yan Liang Fang</i> (Qing Dynasty, AD 1841)

Continued

Preparation Name	Compositions	Clinic uses	References
Danshen Guipi Decoction	<i>Salvia miltiorrhiza</i> Bge., <i>Dipsacus asper</i> Wall. Ex Henry, <i>Paeonia lactiflora</i> Pall., <i>Polygala tenuifolia</i> Willd., <i>Dioscorea opposita</i> Thunb., <i>Fritillaria cirrhosa</i> D.Don, <i>Ophiopogon japonicus</i> (L. f) Ker-Gawl., <i>Poria cocos</i> (Schw.) Wolf, <i>Citrus reticulata</i> Blanco, <i>Nelumbo nucifera</i> Gaertn., <i>Cyathula officinalis</i> Kuan, <i>Rehmannia glutinosa</i> Libosch., <i>Angelica sinensis</i> (Oliv.) Diels, <i>Leonurus japonicus</i> Houtt., <i>Nelumbo nucifera</i> Gaertn.	Treating spitting blood	<i>Chuai Mo You De Ji</i> (Qing Dynasty, AD 1888)
Qingying Decoction	<i>Rehmannia glutinosa</i> Libosch., <i>Cornu Rhinoceri</i> Asiatici, <i>Salvia miltiorrhiza</i> Bge., <i>Scrophularia ningpoensis</i> Hemsl., <i>Ophiopogon japonicus</i> (L.f) Ker-Gawl., <i>Lonicera japonica</i> Thunb., <i>Forsythia suspensa</i> (Thunb.) Vahl, <i>Coptis chinensis</i> Franch., <i>Phyllostachys nigra</i> (Lodd.) Munro var. <i>henanis</i> (Mitf.) Stapf ex Rendle	Treating acute infectious febrile diseases	<i>Wen Bing Tiao Bian</i> (Qing Dynasty, AD 1798)
Qingliang Huagai Yin	<i>Salvia miltiorrhiza</i> Bge., <i>Anemarrhena asphodeloides</i> Bge., <i>Commiphora myrrha</i> Engl., <i>Glycyrrhiza uralensis</i> Fisch.	Treating lung abscess	<i>Yi Xue Zhong Zhong Can Xi Lu</i> (AD 1918–1934)
Huoluo Xiaoling Dan	<i>Salvia miltiorrhiza</i> Bge., <i>Angelica sinensis</i> (Oliv.) Diels, <i>Boswellia carterii</i> Birdw., <i>Commiphora myrrha</i> Engl.	Treating pain in the chest, abdomen, leg, and arm	<i>Yi Xue Zhong Zhong Can Xi Lu</i> (AD 1918–1934)
Erдан Pill	<i>Salvia miltiorrhiza</i> Bge., cinnabar, <i>Polygala tenuifolia</i> Willd., <i>Poria cocos</i> (Schw.) Wolf, <i>Panax ginseng</i> C. A. Mey., <i>Acorus tatarinowii</i> Schott, <i>Rehmannia glutinosa</i> Libosch., <i>Asparagus cochinchinensis</i> (Lour.) Merr., <i>Ophiopogon japonicus</i> (L.f) Ker-Gawl., <i>Glycyrrhiza uralensis</i> Fisch.	Treating amnesia	<i>Bao Ming Ji</i> (Jin Dynasty, AD 1188)
Qingxin Pill	<i>Rehmannia glutinosa</i> Libosch., <i>Salvia miltiorrhiza</i> Bge., <i>Phellodendron chinense</i> Schneid., <i>Ostrea gigas</i> Thunberg, <i>Dioscorea opposita</i> Thunb., <i>Ziziphus jujuba</i> Mill. var. <i>spinosa</i> (Bunge) Hu ex H. F. Chou, <i>Poria cocos</i> (Schw.) Wolf, <i>Ophiopogon japonicus</i> (L.f) Ker-Gawl., <i>Schisandra chinensis</i> (Turcz.) Baill., <i>Plantago asiatica</i> L., <i>Polygala tenuifolia</i> Willd.	Treating seminal emission	<i>Yi Xue Xin Wu</i> (Qing Dynasty, AD 1732)
Danshen Tablet	<i>Salvia miltiorrhiza</i> Bge.	Treating angina pectoris and irritability due to coronary heart disease	<i>Chinese Pharmacopoeia</i>
Fufang Danshen Tablet	<i>Salvia miltiorrhiza</i> Bge., <i>Panax notoginseng</i> (Burk.) F. H. Chen, <i>Cinnamomum camphora</i> (L.) Presl	Treating chest distress and angina pectoris	<i>Chinese Pharmacopoeia</i>
Guanxin Danshen Tablet	<i>Salvia miltiorrhiza</i> Bge., <i>Panax notoginseng</i> (Burk.) F. H. Chen, <i>Dalbergia odorifera</i> T. Chen	Treating chest distress, chest impediment syndrome, palpitation and shortness of breath due to qi stagnation, blood-stasis and coronary heart disease	<i>Chinese Pharmacopoeia</i>
Danqi Tablet	<i>Salvia miltiorrhiza</i> Bge., <i>Panax notoginseng</i> (Burk.) F. H. Chen	Treating impediment and pain of the heart and chest, vertigo, headache and dysmenorrhea due to blood-stasis and qi-stagnation	<i>Chinese Pharmacopoeia</i>
Danyi Tablet	<i>Salvia miltiorrhiza</i> Bge., <i>Leonurus japonicus</i> Houtt., <i>Verbena officinalis</i> L., <i>Achyranthes bidentata</i> Bl., <i>Phellodendron chinense</i> Schneid., <i>Pulsatilla chinensis</i> (Bge.) Regel, <i>Vaccaria segetalis</i> (Neck.) Garcke	Treating stagnation of blood stasis and downward flow of damp-heat	<i>Chinese Pharmacopoeia</i>
Danxi Granule	<i>Salvia miltiorrhiza</i> Bge., <i>Achyranthes bidentata</i> Bl., <i>Gastrodia elata</i> Bl., <i>Paeonia suffruticosa</i> Andr., <i>Paeonia lactiflora</i> Pall., <i>Ligusticum chuanxiong</i> Hort., <i>Rehmannia glutinosa</i> Libosch., <i>Epimedium brevicornu</i> Maxim., <i>Taxillus chinensis</i> (DC.) Danser, <i>Gardenia jasminoides</i> Ellis, <i>Cassia obtusifolia</i> L., <i>Cannabis sativa</i> L.	Treating obstruction of collaterals by blood stasis and deficiency of the kidney due to apoplexy	<i>Chinese Pharmacopoeia</i>
Danxiang Qingzhi Granule	<i>Salvia miltiorrhiza</i> Bge., <i>Ligusticum chuanxiong</i> Hort., <i>Prunus persica</i> (L.) Batsch, <i>Dalbergia odorifera</i> T. Chen, <i>Sparganium stoloniferum</i> Buch.-Ham., <i>Curcuma phaeocaulis</i> Val., <i>Citrus aurantium</i> L., <i>Rheum palmatum</i> L.	Treating qi-stagnation and blood-stasis syndrome due to hyperlipidemia	<i>Chinese Pharmacopoeia</i>
Fufang Danshen Capsule	<i>Salvia miltiorrhiza</i> Bge., <i>Panax notoginseng</i> (Burk.) F. H. Chen, <i>Cinnamomum camphora</i> (L.) Presl	Treating chest impediment syndrome due to qi-stagnation and blood-stasis	<i>Chinese Pharmacopoeia</i>

Continued

Preparation Name	Compositions	Clinic uses	References
Guanxin Danshen Capsule	<i>Salvia miltiorrhiza</i> Bge., <i>Panax notoginseng</i> (Burk.) F. H. Chen, <i>Dalbergia odorifera</i> T. Chen	Treating chest impediment syndrome due to qi-stagnation and blood-stasis	<i>Chinese Pharmacopoeia</i>
Danhong Huayu Oral Liquid	<i>Salvia miltiorrhiza</i> Bge., <i>Angelica sinensis</i> (Oliv.) Diels, <i>Ligusticum chuanxiong</i> Hort., <i>Prunus persica</i> (L.) Batsch, <i>Carthamus tinctorius</i> L., <i>Bupleurum chinense</i> DC., <i>Citrus aurantium</i> L.	Treating obstructive absorption period of central vein of retina attributed to qi-stagnation and blood-stasis syndrome	<i>Chinese Pharmacopoeia</i>
Fufang Danshen Dripping Pill	<i>Salvia miltiorrhiza</i> Bge., <i>Panax notoginseng</i> (Burk.) F. H. Chen, <i>Cinnamomum camphora</i> (L.) Presl	Treating chest impediment syndrome due to qi-stagnation and blood-stasis	<i>Chinese Pharmacopoeia</i>
Fufang Danshen Aerosol	<i>Salvia miltiorrhiza</i> Bge., <i>Panax notoginseng</i> (Burk.) F. H. Chen, <i>Cinnamomum camphora</i> (L.) Presl	Treating chest impediment syndrome due to qi-stagnation and blood-stasis	<i>Chinese Pharmacopoeia</i>

## Chemical Constituents

A total of 201 compounds of Danshen have been summarized so far. These chemical constituents are divided into three groups

based upon their structures: lipophilic diterpenoids, water-soluble phenolic acids and others. The structures of diterpenoids and phenolic acids are shown in Figs. 1–4, and the names of all the constituents are listed in Tables 2–5.







Fig. 1 Structures of tanshinones with a common ortho-naphthoquinone chromophore



**Fig. 2** Structures of abietane diterpenes with a common para-naphthoquinone chromophore



**Fig. 3** Structures of the other components of diterpenoids


**Fig. 4 Structures of phenolic acids**

Strong evidence has proved that diterpenoids and phenolic acids exhibit numerous pharmacological activities, we thereby deduce that they are major representative constituents in Danshen.

#### Diterpenoids

Diterpenoids share a core structure of 20 carbons arranged in rings. These compounds are one group of the major bioactive components in Danshen. This group of compounds possesses a variety of pharmacological activities, such as antibacterial, anti-oxidative, anti-inflammatory, and antineoplastic [2]. At present, at least 81 diterpenoids from Danshen have been reported. According to the structural characteristics, diterpenoids in Danshen are further classified into two sub-sets. One sub-set is tanshinones which are the most abundant diterpenoids, and they contain a common ortho-naphthoquinone chromophore. The other sub-set contains royleanones, which are abietane

diterpenes shared para-naphthoquinone chromophore. Moreover, diterpene chinone compounds of the two sub-sets are found exclusively in the *Salvia* genus [3]. Among all the diterpenoids, tanshinone IIA is the major constituent of the processed herb, and many studies have mainly focused on its pharmacological activities. The biosyntheses of terpenes and terpenoids were studied. It has reported that they could be synthesized through mevalonate (MVA) and methylerythritol phosphate (MEP) pathways [4-6]. The main biosynthetic pathway of tanshinones is MEP pathway, while *SmHMGR1* and *SmHMGR2* from MVA pathway are also beneficial for tanshinone biosyntheses [7-9].

#### Phenolic acids

The water-soluble phenolic acids have been accounted for the therapeutic effects of Danshen. Since 1980s, a lot of Chinese and Japanese researchers have paid attention to these

**Table 2** Diterpenoids isolated from Danshen

No.	Name	References	No.	Name	References
1	11, 12, -dioxoabieta-8, 13-dien (= miltirone)	[188]	42	oleoyl neocryptotanshinone	[189]
2	7 $\beta$ -hydroxy-8, 13-abietadiene-11, 12-dione	[2]	43	miltionone I	[190]
3	4-methylenemiltirone	[2]	44	oleoyl danshenxinkun A	[189]
4	2-isopropyl-8-methylphenanthrene-3, 4-dione	[191]	45	dihydroisotanshinone I	[192]
5	methylenedihydratanshinquinone	[2]	46	isotanshinone I	[193]
6	1, 2, 5, 6-tetrahydratanshinone I	[2]	47	isototanshinone	[194]
7	methylenetanshinquinone	[195]	48	1-ketoisocryptotanshinone	[196]
8	3-hydroxymethylenetanshinquinone	[197]	49	neocryptotanshinone	[198]
9	tanshinone IIA	[192]	50	isotanshinone IIA	[199]
10	tanshindiol A	[200]	51	isotanshinone IIB	[198]
11	tanshindiol B	[200]	52	isocryptotanshinone II	[199]
12	tanshindiol C	[200]	53	danshexinkun C	[201]
13	3 $\alpha$ -hydroxytanshinone IIA	[200]	54	danshexinkun D	[202]
14	tanshinone IIB	[203]	55	sibiriquinone A	[204]
15	methyl tanshinonate	[205]	56	sibiriquinone B	[204]
16	tanshinaldehyde	[205]	57	trijuganone A	[206]
17	nortanshinone	[200]	58	ferruginol	[192]
18	15, 16-dihydratanshinone I	[192]	59	sugiol	[192]
19	tanshinone I	[192]	60	6, 12-dihydroxyabieta-5, 8, 11, 13-tetraen-7-one (= montbretol)	[207]
20	formyltanshinone	[2]	61	11, 12-dihydroxy-20-nor-5(10), 6, 8, 11, 13-abietapentaen-1-one (= arucadiol)	[2]
21	tanshinol A	[208]	62	neotanshinlactone	[209]
22	przewaquinone B	[210]	63	salvianan	[211]
23	cryptotanshinone	[192]	64	neosalvianan	[211]
24	isotanshinone II	[193]	65	salvianen	[211]
25	przewaquinone A	[210]	66	salviadione	[211]
26	hydroxytanshinone IIA	[205]	67	microstegiol	[212]
27	przewaquinone C	[205]	68	epi-cryptoacetalide	[213]
28	methyl dihydronortanshinonate	[214]	69	cryptoacetalide	[213]
29	1, 2-dihydrotanshinquinone	[215]	70	epi-danshenspiroketallactone	[200]
30	1, 2-dihydratanshinquinone	[190]	71	danshenspiroketallactone	[192]
31	dihydronortanshinone	[216]	72	9-isopropyl-2, 2, 5-trimethyl-8H-phenaleno[1, 9bc]furan-8-one (= salvilenone)	[217]
32	1, 2-didehydromiltirone	[2]	73	mltipolone	[218]
33	dihydroisotanshinone II	[194]	74	salviacoccin	[212]
34	1-dehydrotanshinone II A	[207]	75	sclareol	[212]
35	1-dehydromiltirone	[2]	76	salviolone	[218]
36	tanshinone V	[218]	77	tanshinlactone	[2]
37	neo-przewaquinone A	[219]	78	salvilenone	[217]
38	saporthoquinone	[220]	79	miltionone II	[190]
39	neocryptotanshinone II	[207]	80	danshenol A	[200]
40	danshenxinkun B	[192]	81	danshenol B	[208]
41	(-)-danshexinkun A	[221]			

1–38 Tanshinones with a common ortho-naphthoquinone chromophore; 39–57 Abietane diterpenes with a common para-naphthoquinone chromophore; 58–81 The other components of diterpenoids.

**Table 3** Phenolic acids isolated from Danshen

No.	Name	References	No.	Name	References
82	caffeic acid	[16]	104	salvianolic acid N	[212]
83	isoferulic acid	[11]	105	salvianolic acid I	[222]
84	danshensu	[223]	106	salvianolic acid J	[224]
85	3-(3, 4-dihydroxyphenyl)-(2R)-lactamide	[225]	107	salvianolic acid E	[11]
86	salvianolic acid F	[226]	108	salvianolic acid B	[10]
87	salvianic acid C	[16]	109	lithospermic acid B	[11]
88	rosmarinic acid	[227]	110	ethyl lithospermate B	[11]
89	methyl rosmarinate	[16]	111	magnesium lithospermate B	[228]
90	salviaflaside	[229]	112	ammonium-potassium lithospermate B	[230]
91	salvianolic acid D	[11]	113	protocatechuic acid	[16]

Continued

No.	Name	References	No.	Name	References
92	prolithospermic acid	[231]	114	protocatechuic aldehyde	[16]
93	salvianolic acid G	[232]	115	2-(3-methoxy-4-hydroxyphenyl)-5-(3-hydroxypropyl)-7-methoxybenzofuran-3-carbaldehyde	[233]
94	salvinal	[234]	116	ailanthoidol	[233]
95	1-hydroxy-pinoresinol-1-O- $\beta$ -D-glucoside	[234]	117	salvianolic acid L	[235]
96	lithospermic acid	[236]	118	dimethyl salvianolic acid B	[10], 49
97	lithospermic acid monomethyl ester	[194]	119	dimethyl heptamethylsalvianolate B	[10]
98	lithospermic acid dimethyl ester	[16]	120	salvianolic acid K	[237]
99	ethyl lithospermate	[11]	121	ferulic acid	[11]
100	salvianolic acid C	[10]	122	salvianolic acid T	[238]
101	methyl salvianolic acid C	[239]	123	salvianolic acid U	[238]
102	dimethyl lithospermate	[236]	124	salvianolic acid A	[240]
103	9"-methyl lithospermate	[236]			

**Table 4 Essential Oils isolated from Danshen**

No.	Name	References	No.	Name	References
125	borneol acetate	[241]	142	hexadecanoic acid	[241]
126	copaene	[241]	143	linoleic acid	[242]
127	bourbonene	[241]	144	tricosane	[241]
128	iso-elemene	[241]	145	pentacosane	[212]
129	iso- $\beta$ -caryophyllene	[241]	146	heptacosane	[241]
130	isocaryophyllene	[241]	147	nonacosane	[241]
131	$\beta$ -caryophyllene	[241]	148	germacrene B	[212]
132	$\beta$ -cubebene	[241]	149	spathulenol	[212]
133	$\alpha$ -caryophyllene	[241]	150	terpinene-4-ol	[212]
134	cadinadiene	[241]	151	$\alpha$ -Pinene	[212]
135	bicyclogermacrene	[241]	152	$\alpha$ -cadinol	[212]
136	$\alpha$ -farnesene	[241]	153	$\alpha$ -thujene	[212]
137	$\beta$ -caryophyllene oxide	[241]	154	$\beta$ -cadinol	[212]
138	ledol	[241]	155	$\beta$ -chamigrene	[212]
139	$\alpha$ -caryophyllene oxide	[241]	156	$\beta$ -phellandrene	[212]
140	tetradecanoic acid	[241]	157	$\alpha$ -elemene	[212]
141	nor-pristan-2-ol	[241]			

**Table 5 The other components isolated from Danshen**

No.	Name	References	No.	Name	References
158	uvaol	[212]	180	isoleucine	[243]
159	$\alpha$ -amyrin	[212]	181	phenylalanine	[243]
160	$\beta$ -sitosterol- $\beta$ -D-glucoside	[212]	182	valine	[243]
161	luteolin	[212]	183	threonine	[243]
162	baicalin	[212]	184	arginine	[243]
163	daucosterol	[244]	185	serine	[212]
164	$\beta$ -sitosterol	[229]	186	tyrosine	[212]
165	shanzhiside methyl ester	[212]	187	hydrolysis amino acid	[243]
166	manool	[212]	188	calcium	[244]
167	3, 4-dihydroxyphenyl ethanol ketone	[212]	189	cobalt	[245]
168	hexadecane	[212]	190	copper	[245]
169	octadecanol	[212]	191	magnesium	[244]
170	palmitic acid	[242]	192	iron	[244]
171	fructose	[246]	193	nickel	[244]
172	saccharose	[246]	194	zinc	[244]
173	rafitrinose	[246]	195	barium	[244]
174	stachyose	[246]	196	aluminum	[244]
175	vitamin E	[247]	197	stannum	[244]
176	glutamic acid	[243]	198	cadmium	[245]
177	alanine	[243]	199	lead	[245]
178	aspartic acid	[243]	200	manganese	[245]
179	histidine	[243]	201	selenium	[244]

constituents, and more than 20 phenolic acids have been isolated [10–15]. The phenolic acids exhibit broad biological activities, such as anti-oxidative, anti-coagulation, and anti-inflammatory [16–18]. The phenolic acids in Danshen have been classified into single phenolic acids and polyphenolic acids. The single phenolic acids mainly contain a core skeleton of phenylpropanoid (C<sub>6</sub>-C<sub>3</sub>). This group includes danshensu, caffeic acid, protocatechuic aldehyde, protocatechuic acid and many others. Danshensu,  $\beta$ -(3, 4-dihydroxyphenyl) lactic acid, was the first compound that have been discovered among phenolic acids group. The polyphenolic acids are mainly considered as conjugate of Danshensu and derivatives or dimer of caffeic acid. Among which, salvianolic acid A–E, lithospermic acid and rosmarinic acid are depsides. The biosyntheses of phenolic acids in Danshen are involved in phenylpropanoid and tyrosine-derived pathways. It has reported that tyrosine-derived pathway might be the rate-limiting step in the biosynthetic pathways of phenolic acid, because some enzymes of tyrosine-derived pathway were related to the biosyntheses of rosmarinic and salvianolic acid B [19–20].

#### *The other components*

The other components in Danshen include essential oils, triterpenoids, flavone, amino acids, metallic elements, and many others. Among these, 33 essential oils found in Danshen have been concluded in this study, such as borneol acetate, copaene, bourbonene, iso-elemene, iso- $\beta$ -caryophyllene, isocaryophyllene,  $\beta$ -caryophyllene, and  $\alpha$ -caryophyllene. Large number of essential oil constituents have been isolated and identified from the flower of Danshen. In addition, Danshen contains several amino acids, including glutamic acid, alanine, aspartic acid, phenylalanine, serine and many others.

## Quality Control

#### *The application of analytical methods for the quality evaluation*

As is known to all, the quality of Danshen is closely related to the concentrations of their active ingredients. Generally, the water-soluble salvianolic acids and liposoluble tanshinones are considered as the major bioactive components, and hence quality markers in Danshen. Some studies have been conducted to identify the characteristic components in Danshen, and the main bioactive constituents in Danshen have been determined using the three most commonly used analytical methods, including high performance liquid chromatography combined with diode array detector (HPLC-DAD), ultra-high performance liquid chromatography (UPLC), and UPLC coupled with tandem mass spectrometry (UPLC-MS/MS), respectively [21–24]. HPLC-DAD method has been established for respective or simultaneous determination of liposoluble or water-soluble characteristic constituents in Danshen. While UPLC and UPLC-MS/MS methods, which have been considered to be much more rapid and sensitive methods could be employed for the simultaneous quantification of liposoluble and water-soluble constituents in Danshen. Furthermore, some representative researches have utilized UPLC-MS/MS method for comparison and quantification of active compounds in different batches of

Danshen preparations.

#### *The chemical fingerprint analysis*

The chemical fingerprint analysis of traditional Chinese medicine was proposed by the State Food and Drug Administration (SFDA) in 2000. It was accepted for the identification of authenticity, differentiation of origin, and evaluation of quality of herbal medicines and related preparations, which based on the holistic chemical profile obtained by various analytical techniques. There are some reports about fingerprint analysis of Danshen to characterize the whole chemical profile with the assays of HPLC-DAD, UPLC-DAD, and HPLC-MS/MS coupled with chemometrics including hierarchical cluster analysis (HCA), principal component analysis (PCA), and partial least squares discriminant analysis (PLS-DA) [25–26]. The HPLC-DAD and UPLC-DAD methods were developed to observe the multiple common peaks in accordance with different compounds in Danshen. The peaks/compounds in the HPLC or UPLC chromatograms were identified by their characteristic fragment ion information in LC-MS/MS analysis. Furthermore, the characteristic markers and the quality of Danshen were analyzed and evaluated with chemometrics method. And the experimental results suggested varied different bioactive components in Danshen were influenced by processing method to a great degree.

#### *The metabolites analysis in vivo of Danshen decoction or its preparations*

Preclinical researches on the absorption, distribution, metabolism and excretion (ADME) of components in Danshen decoction and its preparations are of great importance for better understanding of their biological effects and safety. Studies on determination of the phenolic constituents of Danshen in serum after intravenous administration, hypodermic administration and oral administration by LC-MS or by microdialysis coupled with HPLC have been reported [27–31].

It has reported that salvianolic acid B has extremely low systemic bioavailability in rats using high performance liquid chromatography with electrochemical detection (HPLC-ECD) method and mostly excreted rapidly into bile as methylated metabolites [32–33]. Moreover, it has indicated that the total phenolic acids or salvianolic acid B are distributed to rat tissues rapidly and the major distribution organ of the four phenolic acids in rat is kidney after oral administration [31].

## Pharmacological Activities

#### *Anti-inflammation*

Inflammatory responses induced by cytokines and chemokines can cause many inflammatory vascular diseases, for which anti-inflammation is a very important therapeutic strategy. Some constituents in Danshen have been investigated for its anti-inflammatory activity [34]. Recent studies have indicated that salvianolic acid B and an aqueous ethanol extract from Danshen possessed anti-inflammatory property. They strongly inhibited tumor necrosis factor- $\alpha$ -induced (TNF- $\alpha$ -induced) nuclear factor- $\kappa$ B (NF- $\kappa$ B) activation in human aortic endothelial cells [35]. Another study has demonstrated

that a novel mechanism for the anti-inflammatory activity of tanshinone IIA may involve down-regulation of vascular cell adhesion molecule-1 and intracellular adhesion molecule-1 through partial blockage of TNF- $\alpha$ -induced NF- $\kappa$ B activation and nuclear factor of kappa light polypeptide gene enhancer in B-cells inhibitor, alpha ( $\text{I}\kappa\text{B}-\alpha$ ) phosphorylation by the inhibition of  $\text{I}\kappa\text{B}$  kinase- $\alpha$  and  $\text{I}\kappa\text{B}$  kinase- $\beta$  pathway in endothelial progenitor cells [36]. Additionally, cryptotanshinone also possessed anti-inflammatory property through various mechanisms, including inhibition of the NF- $\kappa$ B and mitogen-activated protein kinase (MAPK) signaling pathways [37]. Moreover, some constituents isolated from Danshen, although in small quantity, had stronger anti-inflammatory activity than tanshinone IIA, for instance, tanshinone IIB, danshixinkun B, danshenol A, arucadiol, tanshindiol C, salviolone, and sugiol. And it has found that danshenol A ( $5 \mu\text{mol}\cdot\text{L}^{-1}$ ) had the highest inhibition ratio for the secretion of TNF- $\alpha$ , interleukin (IL)-1 $\beta$ , and interleukin (IL)-8 at 56.3%, 67.6%, and 51.7%, respectively [38]. Anti-inflammation contributes significantly in protection against many pathological conditions. Tanshinone IIA has been therapeutically used for the treatment of various diseases, such as acute lung injury, acute kidney injury, fibrosis, Alzheimer's disease (AD) and atherosclerosis, by alleviating inflammation [39-43]. Additionally, it has found that salvianolic acid B could be a potential candidate for the treatment of vascular inflammatory diseases [44-45].

#### Anti-oxidative activity

Oxidative stress is an underlying cause for numerous pathological disorders. Antioxidants may have protective effects against these disorders by keeping reactive species in check. Phenolic acids and tanshinones have antioxidative activity, though the activity of the former is much stronger. Salvianolic acids, especially salvianolic acid A and salvianolic acid B, have been found to have cardiovascular protective effect due to their anti-oxidative activity [46]. Additionally, it has demonstrated that tanshinone IIA contributed to the inhibition of myocardial remodeling, myocardial ischemia reperfusion injury, tanshinone IIA-induced neuroprotection from experimental ischemic stroke, and preventing cirrhosis *via* maintaining antioxidant effect [41, 47-49]. Moreover, owing to antioxidative activity, tanshinone I could provide mitochondrial protection against  $\text{H}_2\text{O}_2$  and protect pyramidal neurons of the gerbil hippocampal CA1 region from ischemic damage induced by transient cerebral ischemia [50-51]. A study has indicated that caffeic acid and its four polymers had the activity of scavenging free radicals. Among which, lithospermic acid B and its  $\text{Mg}^{2+}$  salt displayed the strongest activity ( $\text{IC}_{50}$  of 3.87 and  $4.69 \mu\text{mol}\cdot\text{L}^{-1}$ , respectively). Lithospermic acid and rosmarinic acid also presented good activity ( $\text{IC}_{50}$  of 9.63 and  $10.37 \mu\text{mol}\cdot\text{L}^{-1}$ , respectively), while caffeic acid inhibited 1, 1-diphenyl-2-picrylhydrazyl radical weaker than its polymers, with the  $\text{IC}_{50}$  of  $12.99 \mu\text{mol}\cdot\text{L}^{-1}$  [52].

#### Antitumor activity

Cancer is one of the most common and fatal diseases in the world. Danshen plays an important role in the prevention

of cancer. In one study, Danshen root extract exerted potential cytotoxicity, which can confer sensitivity to various resistant tumors. One possible mechanism is inducing the intrinsic apoptotic pathway [53]. Moreover, several other studies have recently indicated that tanshinones possessed potent anticancer effect both *in vitro* and *in vivo*. It has shown that they suppressed the growth and proliferation of tumor cells, induced apoptosis, inhibited invasion and metastasis, and suppressed angiogenesis *in vitro* [54]. Tanshinone IIA has been found to possess anti-neoplastic activity in the cancer cell-lines of lung, stomach, esophagus, oral cavity, and many others [55-58]. Additional studies have shown that tanshinone IIA induced apoptosis in human leukemia cell lines through the activation of caspase-3 [59] while tanshinone I also played an important role against various tumor cells, such as lung, stomach, breast, and prostate [60-63]. Cryptotanshinone, a potential anticancer agent, could inhibit the mammalian target of rapamycin complex 1 signaling through activation of adenosine monophosphate-activated protein kinase (AMPK)-uberosus sclerosis complex 2 axis in cancer cells [64]. 15, 16-Dihydrotanshinone I could be an efficient therapeutic candidate for osteosarcoma treatment, which is the most common primary malignant bone tumor [65]. Moreover, the antitumor activity of phenolic acids has also been reported, such as salvianolic acid A and salvianolic acid B [66-69]. Salvianolic acid B exerted inhibitory effects on cell proliferation and tumor growth in human glioma U87 cells, which might be related with MAPK 14 p38 activation mediated reactive oxygen species (ROS) generation [70].

#### Effects on cardiovascular and cerebrovascular

##### Anti-atherogenesis

Atherosclerosis is one of the common diseases which are the leading cause of death and disability in our societies. One of the functions of Danshen is removing blood stasis, which indicates its therapeutic value on atherosclerosis treatment and prevention. Many studies have shown that tanshinone IIA had the potential for treating atherosclerosis. Several possible mechanisms have been discovered, such as interfering with receptor of advanced glycation end products and NF- $\kappa$ B activation, inhibition of low density lipoprotein oxidation, anti-inflammation, immunomodulation and many others [43, 71-72]. Cryptotanshinone has been demonstrated to display pleiotropic effects for atherosclerosis through inhibiting lectin-like oxidized low-density lipoprotein receptor-1 (LOX-1) mediated signaling pathway, decreasing ROS level and inhibiting the NF- $\kappa$ B pathways [73-75]. In apolipoprotein E-deficient ( $\text{ApoE}^{-/-}$ ) mice fed with an atherogenic diet, dihydrotanshinone I (10 and  $25 \text{ mg}\cdot\text{kg}^{-1}$ ) significantly attenuated atherosclerotic plaque formation through inhibition of LOX-1 mediated by nicotinamide adenine dinucleotide phosphate-oxidase 4 (NOX4)/NF- $\kappa$ B signaling pathways both *in vitro* and *in vivo* [76]. Moreover, water soluble Danshen extracts showed beneficial effects on atheromatous disease through inhibiting multiple pathways associated with vascular smooth muscle cells, endothelial cells, and platelets both *in vitro* and *in vivo* [77].

### Anti-thrombosis

Thrombosis is hemostasis in the wrong place [78], and the function of Danshen has summarized its anti-thrombotic effect. Danshen is beneficial for the treatment of deep vein thrombosis by its antioxidative effect on vascular endothelial cells [79]. One study has indicated that Danshen exhibited an anti-thrombotic effect, and has clarified its mechanism to be inhibition of the inflammatory response [80]. Experiments have shown that salvianolic acid A reduced platelet adhesion on collagen surfaces by about 40% with an arterial shear rate of  $1000\text{ s}^{-1}$ . Salvianolic acid A could attenuate platelet activation and arterial thrombus formation through inhibiting phosphoinositide 3-kinase (PI3K) [81]. Another study has demonstrated that protocatechuic aldehyde possesses the strong anti-thrombotic activity associated with suppressing the platelet aggregation, using an impedance aggregometer [82]. Moreover, rosmarinic acid could inhibit the formation of venous thrombosis and platelet aggregation by 54.8% and 46.4% at a dosage of  $100\text{ mg}\cdot\text{kg}^{-1}$  [16].

### Anti-hypertension

System hypertension is an asymptomatic disorder, which poses a risk of coronary thrombosis, stroke, and renal failure [83]. Danshen has remarkable beneficiary effects on the treatment of hypertension. Experimental studies have indicated that Danshen may be useful for monocrotaline-induced pulmonary hypertension rats at low or high oral dose ( $4.6$  or  $14\text{ g}\cdot\text{kg}^{-1}$ ) for 21 days [84]. Additionally, water extract of Danshen exerted anti-hypertensive effect by inhibiting the angiotensin converting enzyme activities in the renin-angiotensin system [85]. It has also demonstrated that optimal compatibility ratio of active ingredients from Danshen, including salvianolic acid A, salvianolic acid B, danshensu and protocatechuic aldehyde, exhibited a crucial anti-hypertensive effect on rats. Its mechanism is inhibiting oxidative stress and the transforming growth factor-beta (TGF- $\beta$ )/Smad pathway [86]. One study has shown that sodium tanshinone IIA sulfonate had therapeutic value on pulmonary hypertension patients both alone as well as in concert with sildenafil [87]. Endothelial dysfunction could result in hypertension. It has found that salvianolic acid B had anti-hypertensive effect through reversing the impaired endothelial function and inhibiting  $\text{AT}_1$  receptor-dependent vascular oxidative stress [88].

### Anti-hyperlipidemic

Dyslipidaemia is another risk factor for atheroma and relative cardiovascular diseases and extracts of Danshen demonstrated their effects on lipid regulation in various studies [83]. It has indicated that Danhong injection could adjust lipid metabolic disorders, for example, falling serum lipid levels and suppressing formation of lipid peroxidation products [89]. Moreover, Fufang Danshen Dripping pill could decrease serum lipid levels and protect liver function, which might be associated with ameliorating of anti-oxidation and falling of inflammation [90]. Tanshinone IIA has been found to decrease the lipid deposition in liver. Additionally, it could decrease high density lipoprotein (HDL) middle subfractions

levels and increase HDL large subfractions levels, which regulated intake and efflux of cholesterol [91]. It has shown that cryptotanshinone had anti-adipogenic activity through regulating signal transducer and activator of transcription 3 during early adipogenesis [92]. Moreover, salvianolic acid B and paeonol ( $5$ ,  $10$ , and  $15\text{ mg}\cdot\text{kg}^{-1}\text{ BW}$ ) could markedly and dose-dependently increase HDL levels [93].

### Effects on ischaemic stroke

Thrombosis of a major cerebral artery is a major cause of ischemic brain damage. Danshen and its extracts provide a potential therapeutic approach for ischaemic stroke. It has demonstrated that Danshen dripping pill might be beneficial to treat stroke/transient ischemic attack recurrence *via* anti-inflammation [94]. It has indicated that salvianolic acid B had a potential effect of treating/alleviating brain injury by activating silent information regulator 1 signaling, which was related with ischaemic stroke [95]. Magnesium lithospermate B could also provide neuroprotective effect against ischemic stroke [96]. Additionally, tanshinone IIA has been found to have neuroprotection on experimental ischemic stroke, which may be associated with mechanisms of anti-inflammatory, anti-oxidative, anti-apoptosis, and inhibition of excitatory amino acid toxicity [97]. Tanshinone I and cryptotanshinone also displayed a protective effect against stroke, and the disease status could increase the brain access of cryptotanshinone [51, 98].

### Effects on the heart

A recent study has indicated that Danshen plays an important role in therapeutic interventions for patients with coronary heart disease. It may decrease the risk of coronary heart disease with improved biomarkers of patients [99]. It has demonstrated that aspirin in combination with Fufang Danshen Dripping pill is much more effective than aspirin as monotherapy for treating coronary heart [100]. Myocardial ischemia-reperfusion injury (MIRI) is unavoidable during cardioplegic arrest and open-heart surgery. Danshen can protect against MIRI through various mechanisms. For example, tanshinone IIA can protect against MIRI by activating the PI3K/protein kinase B (Akt)/mTOR signaling pathway, reducing oxidative stress, decreasing HMGB1 expression, and suppressing inflammatory response [48, 101-103]. Dihydro-tanshinone I possessed cardio-protective effect against MIRI through inhibiting arachidonic acid  $\omega$ -hydroxylase [104]. Additionally, Danshensu showed a cardioprotective effect on isolated heart of rats via activation of Akt/extracellular regulated protein kinases (ERK1/2)/nuclear factor erythroid-2-related factor 2 (Nrf2) signaling [105]. Tanshinone IIA can also inhibit myocardial remodeling induced by pressure overload, reduce atrial fibrillation in chronic heart failure and attenuate cardiac dysfunction in endotoxin-induced septic mice [47, 106-107].

### Effects on nervous system

#### Sedative and analgesic activities

Danshen has been widely used for the treatment of neuroathenic insomnia in China. A number of pharmacological studies have been done on the neuroprotection of Danshen. It has demonstrated that miltirone has sedative and muscle re-

laxation activity, and it is showed to be effective in behavioral tests [108]. One study has indicated that the combination of ether extract of Danshen (300 and 600 mg·kg<sup>-1</sup> body wt.) and the water extract of Suanzaoren (400 and 800 mg·kg<sup>-1</sup> body wt.) have a significant sedative-hypnotic activity, which can decrease sleep latency and increase sleeping time induced by sodium pentobarbital (55 mg·kg<sup>-1</sup> body wt.) in mice [109]. Experiments using extracellular microelectrode method and stereotaxic technique of brain on visceral pain discharges on the posterior nucleus of thalamus in cats have shown that Danshen exerted analgesic effect on central nervous system [110]. Evidences have demonstrated that tanshinone IIA relieved spinal nerve ligation-induced neuropathic pain *via* depressing microglial activation and immune response [111]. Moreover, it has found that tanshinone IIA sulfonate has a strong analgesic activity on spinal nerve ligation-induced neuropathic pain by suppressing astrocyte-related c-Jun N-terminal kinase phosphorylation and monocyte chemoattractant protein-1 [112].

#### *Effects on neurodegenerative disorder*

AD is a common neurodegenerative disorder that severely affects millions of elderly people and brings a great social burden. Danshen with protective effects on neurons can be used to treat AD. Tanshinone IIA has been identified to be an effective agent for AD therapy, which can suppress the proliferation of astrocytes, up-regulate the expression of Akt, and inhibit the production of NF- $\kappa$ B and caspase-3 in an AD model [113-114]. Moreover, tanshinone IIA can inhibit inducible nitric oxide synthase (iNOS), matrix metalloproteinase-2 production, and nuclear transcription factor kappa transcription and translation in the temporal lobes of AD rat models, thus exerting neuroprotective effects [115]. Cryptotanshinone also showed a promising potential in the treatment or prevention of AD [98, 116-119]. In one study, cryptotanshinone can inhibit human acetylcholinesterase and butyrylcholinesterase with IC<sub>50</sub> of 4.09 and 6.38  $\mu$ mol·L<sup>-1</sup> in AD model [118]. Dihydrotanshinone has been found to have anti-cholinesterase activity, which indicated it is potential for treating AD [119]. Additionally, phenolic acids were also accounted for the beneficial effects of Danshen in AD treatment, such as salvianolic acid A, salvianolic acid B, and danshensu [120-124]. Salvianolic acid B has been found to inhibit amyloid- $\beta$  generation *via* regulating  $\beta$ -secretase 1 activity in SH-SY5Y-APPsw cells [123].

Parkinson's disease (PD) is a chronic progressive degenerative disorder of the central nervous system, which has the highest incidence in the elderly, especially people over 50 [125]. In one study, tanshinone I and tanshinone IIA can inhibit the aggregation of  $\alpha$ -synuclein both *in vitro* and in a transgenic *Caenorhabditis elegans* PD model [126]. Another experiment has displayed that tanshinone I can prevent nigrostriatal dopaminergic neurodegeneration in a mouse PD model [127]. Oxidative stress caused by dopamine may play an important role in the PD pathogenesis. Tanshinone I can upregulate nuclear factor erythroid-2-related factor 2 to attenuate 6-hydroxydopamine-induced oxidative stress in cellular and mouse PD model [128]. Tanshinone IIA can also prevent the

loss of nigrostriatal dopaminergic neurons *via* the inhibition of nicotinamide adenine dinucleotide phosphate (NADPH) oxidase and iNOS in a 1-methyl-4-phenyl-1, 2, 3, 6-tetrahydropyridine mouse PD model [129]. Moreover, phenolic acids, especially salvianolic acid A, salvianolic acid B and danshensu, were effective in treating PD associated with oxidative stress [124, 130-131]. It has found that salvianolic acid B can protect dopaminergic neurons by an Nrf2-mediated dual action in PD models [125].

#### *Anti-fibrotic activity*

##### *Effects on hepatic fibrosis*

Danshen is used to treat hepatocyte injury and hepatic fibrosis, for it can increase the regeneration of hepatocytes and improve hepatic blood circulation. And it has found that Danshen possessed hepatoprotective effects in chronic iron-overloaded mice [132]. Immunohistochemical examinations have indicated that the hot-water extract of Danshen effectively ameliorated hepatic fibrosis induced by biliary obstruction in rats [133]. Danshen extracts can prevent dimethylnitrosamine-induced hepatic fibrosis by downregulating TGF- $\beta$ 1 [134]. A study has demonstrated that tanshinone IIA can suppress fibrosis and reduce liver injury in a rat model of cirrhosis *via* regulating heme oxygenase-1, Akt and phosphorylated-p38 MAPK signaling pathway [41]. Tanshinone IIA can also protect the liver from acetaminophen-induced hepatic injury through activating Nrf2 pathway [135]. Cryptotanshinone, dihydrotanshinone I and protocatechuic aldehyde also had hepatoprotective effect [136-138]. Salvianolic acid A, salvianolic acid B, magnesium lithospermate B and lithospermic acid played an important role in protecting liver [139-142].

##### *Effects on pulmonary fibrosis*

In addition to hepatic fibrosis, Danshen and its extracts are effective against fibrotic lung diseases as well. One experimental study has shown that Danshen had therapeutic potential in treating fibrotic lung diseases [143-144]. Another experimental study has demonstrated that tanshinone IIA exerted protective effect against bleomycin-induced pulmonary fibrosis in rats [145]. Salvianolic acid A was responsible for preventing pulmonary fibrosis through inhibiting proliferation, adhesion and migration of fibroblasts, and promoting apoptosis [146]. Additionally, salvianolic acid B possessed dose-dependent inhibitory effect on TGF- $\beta$ 1-induced proliferation and differentiation in treating pulmonary fibrosis [147]. IH764-3, a potent component isolated from Danshen, can also be considered as a potential prophylactic and therapeutic agent for fibrotic lung diseases [148-149].

##### *Protective effects against renal injury*

Danshen had protective effects on renal injury induced by myocardial infarction *via* restoring the renal function and preventing the oxidative stress [150]. The ethanol extracts and water extracts of Danshen had therapeutic values on treating renal injury [151-152]. Tanshinone IIA could protect against folic acid-induced kidney injury by attenuating renal tubular epithelial injury and inhibiting local inflammatory response [40]. It could prevent the progression of chronic kidney disease after acute kidney injury *via* fibrocytes recruitment in a mouse

model [153]. Tanshinone IIA could also prevent transition of acute kidney injury to chronic kidney disease by targeting glycogen synthase kinase 3 $\beta$  [154]. Moreover, tanshinone IIA pretreatment could attenuate ischemia/reperfusion-induced renal injury through the inhibition of myeloperoxidase, macrophage migration inhibitory factor, cleaved caspase-3 and p38 MAPK [155]. It has demonstrated that lithospermic acid B could improve ischemia/reperfusion-induced renal injury in rats at a dosage of 40 mg·kg<sup>-1</sup>·day<sup>-1</sup> for 4 days, the mechanism may involve scavenging of reactive oxygen species [156]. Magnesium lithospermate B with therapeutic values on ameliorating the development of age-related renal damage was isolated from the herb, and it promoted restore of renal function through NADPH oxidase-mediated reactive oxygen generation [157]. Additionally, Danshen injection was widely used to ameliorate renal damage induced by iron overload and lead exposure in mice [158-159].

#### Protective effects against diabetes mellitus

Diabetes mellitus, a major endocrine disorder, is an increasingly common disease in many developed and developing countries. Danshen has been showed to exert therapeutic potential against this disorder. One study has indicated that the total polyphenolic acids fraction of Danshen displayed positive effects on type 2 diabetes mellitus rat model at an oral dose (187 mg·kg<sup>-1</sup>) for 28 days [160]. Salvianolic acid A could ameliorate diabetic foot problems through protecting diabetic rats from neuropathy and peripheral vascular circulation dysfunction [161]. Salvianolic acid B exhibited antidiabetic activity through many underlying mechanisms, such as attenuation of oxidative stress, prevention of apoptosis, modulation of the AMPK pathway, suppression of endothelial cell apoptosis and stimulation of endothelial nitric oxide synthase phosphorylation [162-164]. Lithospermic acid B has been reported to exert antioxidant and anti-inflammatory effects, which could treat diabetic retinopathy in rats at a dosage of 10 or 20 mg·kg<sup>-1</sup>·day<sup>-1</sup> for 52 weeks [165]. It has found that danshensu could produce significant protective effects in diabetes [166-167]. Additionally, pioglitazone, an extract of Danshen dripping pill, played an important role in ameliorating diabetic nephropathy [168]. Moreover, Danshen injection was also widely used to treat type 2 diabetes mellitus [169-171], in which the main active components are salvianolic acid A, salvianolic acid B, danshensu, rosmarinic acid and lithospermate B [172-173].

#### Other effects

Two hydrophilic compounds isolated from Danshen, have been discovered to be effective against human immunodeficiency virus type 1 (HIV-1) integrase with no toxic side effects, which made them extremely appealing as a potential of therapeutic drugs for acquired immunodeficiency syndrome [174]. It also has demonstrated that tanshinone II A could inhibit Tat-induced HIV-1 transactivation *via* redox signaling pathway [175]. Besides, Danshen could enhance the gastric mucosal barrier and promote the gastric mucosal cell proliferation along the edge of the ulcer, which played an important role in promoting ulcer healing and preventing

recurrence [176]. Salvianolic acid A exerted anti-secretory and antiulcer effects *via* inhibiting pig gastric H<sup>+</sup>, K<sup>+</sup>-ATPase and pNPPase with IC<sub>50</sub> of 5.2 × 10<sup>-7</sup> mol·L<sup>-1</sup> and 1.7 × 10<sup>-6</sup> mol·L<sup>-1</sup> [177]. Some active components from Danshen, especially 15, 16-dihydrotanshinone I and cryptotanshinone, have been demonstrated to have anti-allergic activity *in vitro* through inhibiting the release of  $\beta$ -hexosaminidase [178-179]. In addition, ethanol extract from Danshen has been found to be responsible for the beneficial effects in the treatment of allergy-related disorders [180]. Danshen has been reported to produce a protective effect against psoriasis [181]. Moreover, cryptotanshinone and dihydrotanshinone I showed antibacterial activity against a broad range of Gram positive bacteria, the mechanism may be related to the generation of superoxide radicals in *Bacillus subtilis* lysate [182]. Salvianolate displayed potentiating activity of multiple antibacterial agents against methicillin-resistant *Staphylococcus aureus in vitro*, which would be beneficial to combinatory therapy for fighting with this infectious pathogen [183].

#### Side Effects

Danshen has been successfully used in the treatment of pregnancy-induced hypertension by inhibiting the angiotensin-converting enzyme. However, many researches have demonstrated that the use of angiotensin-converting enzyme inhibitors was closely related to fetal toxicity, even stillbirths, which were caused by the chemical components from Danshen in the second and third trimesters of pregnancy [184]. It has shown that deposite salt injection made from Danshen had some adverse drug events, such as headache, facial flushing, dizziness skin itching, thrombocytopenia, and abnormal liver function, which may be caused by rapid infusion rate and others. Long-term toxicity tests on beagles showed the safety dose of deposite salt injection of Danshen was lower than 80 mg·kg<sup>-1</sup>, and a dose at 320 mg·kg<sup>-1</sup> showed toxicity [185-186]. One study has indicated that high-dose (5.76 g·kg<sup>-1</sup>·day<sup>-1</sup>) Danshen injection could result in peripheral vascular toxicities, for instance, the increase of vascular leakage, the rise of serum nitrate and endothelin-1, and the apoptosis of vascular endothelial cells [187]. In addition, another research has evaluated the safety of Danshen and Fufang Danshen injection with 2715 patients from 35 randomized controlled trials. Among these, five trials discovered some minor adverse drug events, such as stomach discomfort, itchy skin and local pain, which were reported to be tolerable [185].

At present, no severe adverse drug event on Danshen have been reported, and Danshen products are widely used on its recommended indications at its clinical dose. In order to conduct healthcare services safely and efficiently, more therapeutic evidence on the safety and efficacy of Danshen and its products should be collected.

#### Discussions and Conclusions

This review summarized the latest advancements of Danshen in traditional uses, phytochemistry, quality control, pharmacological activities, and side effects. As a Chinese herbal medicine,

Danshen has been used in therapeutic for various syndromes and preventive measures for thousands of years. Nevertheless, we proposed several suggestive points attempted to enhance the beneficial effects of Danshen in healthcare.

A total of reported 201 chemical constituents of Danshen were summarized in the present review, among which 81 were liposoluble compounds, 43 were water-soluble and 77 were catalogued as “other” constituents. The liposoluble constituents, represented by tanshinones, possessed prominent effects, such as anti-inflammation, anti-cancer, and antibacterial [2]. The water-soluble constituents, represented by phenolic acids, exhibited pharmacological activities including anti-oxidative, anti-coagulation, and anti-atherogenesis [16-18, 71]. Danshen, displaying wide-spectrum pharmacological activities, has been used to alleviate hypertension, dyslipidaemia, diabetes, and others.

As a traditional herbal medicine, Danshen has been commonly used in treating chest impediment, irregular menstruation, stroke, and so on. Among these, chest impediment is a syndrome related to cardiovascular diseases, such as atherosclerosis, thrombosis, angina and coronary heart disease. As is well known, Fufang Danshen Dripping pill, used to treat cardiovascular diseases, was the first Chinese patent traditional medicine to pass Phase II trials of the US Food and Drug Administration. Now, with the developments of medicine and clinical study, the clinical application of Danshen has broadened. As a result, it has shown that Danshen exerted protective effects against diabetes mellitus and cancer.

Diabetes is a chronic disease increasing in prevalence throughout the world. The disease, which is difficult to treat, is usually accompanied with many complications. Herbal medicines show outstanding effects in diabetic therapy. Among which, Danshen has been found to possess antidiabetic activity in clinical trials, and it can effectively be used to treating several diabetic complications in humans. Phenolic acids, such as salvianolic acid A, salvianolic acid B, lithospermic acid B, rosmarinic acid and danshensu, displayed statistically valid results in alleviating diabetes. Additionally, the preparations of Danshen, including Fufang Danshen Dripping pill and Danshen injection, have been used in ameliorating diabetes, and have shown promising outcomes.

Cancer is always one of the most common and refractory diseases that threaten people's health and life. New therapeutic strategies are required for the treatment of numerous cancers. As a widely investigated herb worldwide, the highly medicinal properties of Danshen have gotten increasing attention. Various compounds isolated from Danshen, particularly tanshinones, have been found to exert potent anticancer activity *via* promoting apoptotic cell death and cell cycle arrest and many other mechanisms in cancer cells. Moreover, salvianolic acid A and salvianolic acid B have been reported for antitumor activity.

With increasing formulated herbal products containing Danshen and an expanding global market of those remedies, the quality control of Danshen met new challenges. First of all, different resources and harvest time vary the active compounds

content in Danshen. And the activity of the ingredients is greatly influenced by processing methods, formulation and manufacture processes. To ensure end-product quality, the quality of the processed herb should be strictly controlled. The summarized components could be used to enrich the chemical fingerprint data of Danshen. This review provided up-to-date information on the active components in Danshen and their corresponding pharmacological activity. Such information can be used to compose and improve the quality control standard on Danshen.

Despite that lipophilic and water-soluble constituents in Danshen have been used for the complex biological activities they displayed, the mechanism of action of those compounds remained unclear and required further inspection. A great deal of work should be carried out in order to thoroughly clarify the relationship between compounds and activities. In addition, scientists still need to improve the systematic phytochemistry researching method through analysis. With the development of modern analytical methods, UPLC-MS/MS has been widely used for detection of various constituents. And with the accumulation of analytical data, scientists will isolate new compounds more easily. It is also important to further research some constituents, which have not been tested biologically. Scientists need to work hard to make products containing Danshen be widely applied to clinical practice, which will be advantageous for a holistic assessment of Danshen.

In summary, Danshen, as a folk and traditional herbal medicine, has been widely used in China and other Asian countries. The traditional uses, phytochemical information, pharmacological studies and side effects should be comprehensively understood for its clinical use. Researches on the mechanisms and quality control need to be explored in order to improve its uses in the near future.

## References

- [1] Zhou LM, Zuo Z, Chow MSS. Danshen: an overview of its chemistry, pharmacology, pharmacokinetics, and clinical use [J]. *J Clin Pharmacol*, 2005, **45**(12): 1345-1359.
- [2] Chang HM, Cheng KP, Choang TF, et al. Structure elucidation and total synthesis of new tanshinones isolated from *Salvia miltiorrhiza* Bunge (Danshen) [J]. *J Org Chem*, 1990, **55**(11): 3537-3543.
- [3] Ryu SY, Lee CO, Choi SU. *In vitro* cytotoxicity of tanshinones from *Salvia miltiorrhiza* [J]. *Planta Med*, 1997, **63**(4): 339-342.
- [4] Lichtenthaler HK. Non-mevalonate isoprenoid biosynthesis: enzymes, genes and inhibitors [J]. *Biochem Soc Trans*, 2000, **28**(6): 785-789.
- [5] Laule O, Fürholz A, Chang HS, et al. Crosstalk between cytosolic and plastidial pathways of isoprenoid biosynthesis in *Arabidopsis thaliana* [J]. *Proc Natl Acad Sci USA*, 2003, **100**(11): 6866-6871.
- [6] Ge XC, Wu JY. Tanshinone production and isoprenoid pathways in *Salvia miltiorrhiza* hairy roots induced by Ag<sup>+</sup> and yeast elicitor [J]. *Plant Sci*, 2005, **168**(2): 487-491.

- [7] Kai GY, Xu H, Zhou CC, et al. Metabolic engineering tanshinone biosynthetic pathway in *Salvia miltiorrhiza* hairy root cultures [J]. *Metab Eng*, 2011, **13**(3): 319-327.
- [8] Shi M, Lou X, Ju G, et al. Increased accumulation of the cardio-cerebrovascular disease treatment drug tanshinone in *Salvia miltiorrhiza* hairy roots by the enzymes 3-hydroxy-3-methylglutaryl CoA reductase and 1-deoxy-D-xylulose 5-phosphate reductoisomerase [J]. *Funct Integr Genomics*, 2014, **14**(3): 603-615.
- [9] Dai ZB, Cui GH, Zhou SF, et al. Cloning and characterization of a novel 3-hydroxy-3-methylglutaryl coenzyme A reductase gene from *Salvia miltiorrhiza* involved in diterpenoid tanshinone accumulation [J]. *J Plant Physiol*, 2011, **168**(2): 148-157.
- [10] Ai CB, Li LN. Stereostructure of salvianolic acid B and isolation of salvianolic acid C from *Salvia miltiorrhiza* [J]. *J Nat Prod*, 1988, **51**(1): 145-149.
- [11] Ai CB, Li LN. Salvianolic acids D and E: two new depsides from *Salvia miltiorrhiza* [J]. *Planta Med*, 1992, **58**(2): 197-199.
- [12] Huang L, Liu D, Hu Z. Effects of phytohormones on growth and content of depsides in *Salvia miltiorrhiza* suspension cells [J]. *Chin Med Mat*, 2000, **23**(1): 1-4.
- [13] Li J, He LY, Song WZ. Separation and quantitative determination of seven aqueous depsides in *Salvia miltiorrhiza* by HPTLC scanning [J]. *Acta Pharm Sin*, 1993, **28**(7): 543-547.
- [14] Li LN, Tan R, Chen WM. Salvianolic acid A, a new depside from roots of *Salvia miltiorrhiza* [J]. *Planta Med*, 1984, **50**(3): 227-228.
- [15] Murakami S, Kijima H, Isobe Y, et al. Effect of salvianolic acid A, a depside from roots of *Salvia miltiorrhiza*, on gastric H<sup>+</sup>, K<sup>+</sup>-ATPase [J]. *Planta Med*, 1990, **56**(4): 360-363.
- [16] Jiang RW, Lau KM, Hon PM, et al. Chemistry and biological activities of caffeic acid derivatives from *Salvia miltiorrhiza* [J]. *Curr Med Chem*, 2005, **12**(2): 237-246.
- [17] Du GH, Zhang JT. The general situation and progress of the modern research of red sage root (*Radix Salviae miltiorrhizae*). I. [J]. *Herald Med*, 2004, **23**: 355-360.
- [18] Du GH, Zhang JT. The general situation and progress of the modern research of red sage root (*Radix Salviae miltiorrhizae*). II. [J]. *Herald Med*, 2004, **23**: 435-440.
- [19] Zhang SC, Yan Y, Wang BQ, et al. Selective responses of enzymes in the two parallel pathways of rosmarinic acid biosynthetic pathway to elicitors in *Salvia miltiorrhiza* hairy root cultures [J]. *J Biosci Bioeng*, 2014, **117**(5): 645-651.
- [20] Yan Q, Shi M, Ng J, et al. Elicitor-induced rosmarinic acid accumulation and secondary metabolism enzyme activities in *Salvia miltiorrhiza* hairy roots [J]. *Plant Sci*, 2006, **170**(4): 853-858.
- [21] Chen TL, Bi CQ, Xiao X, et al. Fingerprint and simultaneous determination of multi-components in water-soluble components of *Salvia miltiorrhiza* in Miao Autonomous County of Songtao, Guizhou [J]. *Chin Med Mat*, 2015, **38**(3): 536-539.
- [22] Liu AH, Lin YH, Yang M, et al. Development of the fingerprints for the quality of the roots of *Salvia miltiorrhiza* and its related preparations by HPLC-DAD and LC-MS(n) [J]. *J Chromatogr B Analyt Technol Biomed Life Sci*, 2007, **846**(1-2): 32-41.
- [23] Zhang LL, Liu YY, Liu ZL, et al. Comparison of the roots of *Salvia miltiorrhiza* Bunge (Danshen) and its variety *S. miltiorrhiza* Bge f. Alba (Baihua Danshen) based on multi-wavelength HPLC-fingerprinting and contents of nine active components [J]. *Anal Methods-UK*, 2016, **8**(15): 3171-3182.
- [24] Luo H, KW, Hu Y, et al. Quality evaluation of *Salvia miltiorrhiza* Bge. by ultra high performance liquid chromatography with photo-diode array detection and chemical fingerprinting coupled with chemometric analysis [J]. *J Sep Sci*, 2015, **38**(9): 1544-1551.
- [25] Liu M, Li YG, Chou GX, et al. Extraction and ultra-performance liquid chromatography of hydrophilic and lipophilic bioactive components in a Chinese herb *Radix Salviae Miltiorrhizae* [J]. *J Chromatogr A*, 2007, **1157**(1-2): 51-55.
- [26] Liang WY, Chen WJ, Wu LF, et al. Quality evaluation and chemical markers screening of *Salvia miltiorrhiza* Bge. (Danshen) based on HPLC fingerprints and HPLC-MS(n) coupled with chemometrics [J]. *Molecules*, 2017, **22**(3): E478.
- [27] Li XC, Yu C, Sun WK, et al. Pharmacokinetics of magnesium lithospermate B after intravenous administration in beagle dogs [J]. *Acta Pharmacol Sin*, 2004, **25**(11): 1402-1407.
- [28] Li XC, Yu C, Sun WK, et al. Simultaneous determination of magnesium lithospermate B, rosmarinic acid, and lithospermic acid in beagle dog serum by liquid chromatography/tandem mass spectrometry [J]. *R Rapid Commun Mass Spectrom*, 2004, **18**(23): 2878-2882.
- [29] Li X, Yu C, Sun W, et al. Liquid chromatography/tandem mass spectrometry for the determination of magnesium lithospermate B in beagle dog serum [J]. *J Pharm Biomed Anal*, 2005, **38**(1): 107-111.
- [30] Li X, Yu C, Cai Y, et al. Simultaneous determination of six phenolic constituents of danshen in human serum using liquid chromatography/tandem mass spectrometry [J]. *J Chromatogr B Analyt Technol Biomed Life Sci*, 2005, **820**(1): 41-47.
- [31] Xu M, Fu G, Qiao X, et al. HPLC method for comparative study on tissue distribution in rat after oral administration of salvianolic acid B and phenolic acids from *Salvia miltiorrhiza* [J]. *Biomed Chromatogr*, 2007, **21**(10): 1052-1063.
- [32] Zhang Y, Akao T, Nakamura N, et al. Extremely low bioavailability of magnesium lithospermate B, an active component from *Salvia miltiorrhiza*, in rat [J]. *Planta Med*, 2004, **70**(2): 138-142.
- [33] Zhang Y, Akao T, Nakamura N, et al. Magnesium lithospermate B is excreted rapidly into rat bile mostly as methylated metabolites, which are potent antioxidants [J]. *Drug Metab Dispos*, 2004, **32**(7): 752-757.
- [34] Zhou X, Razmovski-Naumovski V, Chang D, et al. Synergistic effects of Danshen (*Salvia miltiorrhiza* Radix et Rhizoma) and Sanqi (Notoginseng Radix et Rhizoma) combination in inhibiting inflammation mediators in RAW264.7 cells [J]. *Biomed Res Int*, 2016, **2016**:1-12.
- [35] Chen YH, Lin SJ, Ku HH, et al. Salvianolic acid B attenuates VCAM-1 and ICAM-1 expression in TNF-alpha-treated human aortic endothelial cells [J]. *J Cell Biochem*, 2001, **82**(3): 512-521.

- [36] Yang JX, Pan YY, Ge JH, et al. Tanshinone II A attenuates TNF- $\alpha$ -induced expression of VCAM-1 and ICAM-1 in endothelial progenitor cells by blocking activation of NF- $\kappa$ B [J]. *Cell Physiol Biochem*, 2016, **40**(1-2): 195-206.
- [37] Tang S, Shen XY, Huang HQ, et al. Cryptotanshinone suppressed inflammatory cytokines secretion in RAW264.7 macrophages through inhibition of the NF- $\kappa$ B and MAPK signaling pathways [J]. *Inflammation*, 2011, **34**(2): 111-118.
- [38] Ma SL, Zhang DW, Lou HX, et al. Evaluation of the anti-inflammatory activities of tanshinones isolated from *Salvia miltiorrhiza* var. *alba* roots in THP-1 macrophages [J]. *J Ethnopharmacol*, 2016, **188**: 193-199.
- [39] Xu M, Cao FL, Zhang YF, et al. Tanshinone IIA therapeutically reduces LPS-induced acute lung injury by inhibiting inflammation and apoptosis in mice [J]. *Acta Pharmacol Sin*, 2015, **36**(2): 179-187.
- [40] Jiang CM, Zhu W, Shao QY, et al. Tanshinone IIA protects against folic acid-induced acute kidney injury [J]. *Am J Chin Med*, 2016, **44**(4): 737-753.
- [41] Shu M, Hu XR, Hung ZA, et al. Effects of tanshinone IIA on fibrosis in a rat model of cirrhosis through heme oxygenase-1, inflammation, oxidative stress and apoptosis [J]. *Mol Med Rep*, 2016, **13**(4): 3036-3042.
- [42] Lu BL, Li J, Zhou J, et al. Tanshinone IIA decreases the levels of inflammation induced by A $\beta$ 1-42 in brain tissues of Alzheimer's disease model rats [J]. *Neuroreport*, 2016, **27**(12): 883-893.
- [43] Zhao D, Tong L, Zhang L, et al. Tanshinone II A stabilizes vulnerable plaques by suppressing RAGE signaling and NF- $\kappa$ B activation in apolipoprotein-E-deficient mice [J]. *Mol Med Rep*, 2016, **14**(6): 4983-4990.
- [44] Xu SX, Zhong AQ, Bu XK, et al. Salvianolic acid B inhibits platelets-mediated inflammatory response in vascular endothelial cells [J]. *Thromb Res*, 2015, **135**(1): 137-145.
- [45] Xie LX, Durairajan SSK, Lu JH, et al. The effect of salvianolic acid B combined with laminar shear stress on TNF- $\alpha$ -stimulated adhesion molecule expression in human aortic endothelial cells [J]. *Clin Hemorheol Microcirc*, 2010, **44**(4): 245-258.
- [46] Ho JHC, Hong CY. Salvianolic acids: small compounds with multiple mechanisms for cardiovascular protection [J]. *J Biomed Sci*, 2011, **18**(1): 18-30.
- [47] Feng J, Li SS, Chen HW. Tanshinone IIA inhibits myocardial remodeling induced by pressure overload via suppressing oxidative stress and inflammation: possible role of silent information regulator 1 [J]. *Eur J Pharmacol*, 2016, **791**: 632-639.
- [48] Hu HL, Zhai CL, Qian G, et al. Protective effects of tanshinone IIA on myocardial ischemia reperfusion injury by reducing oxidative stress, HMGB1 expression, and inflammatory reaction [J]. *Pharm Biol*, 2015, **53**(12): 1752-1758.
- [49] Cai M, Guo Y, Wang S, et al. Tanshinone II A elicits neuroprotective effect via activating the nuclear factor erythroid 2-related factor -dependent antioxidant response [J]. *Rejuv Res*, 2017, **20**(4): 286-297.
- [50] Oliveira MR, Fürstenau CR, Ferreira GD, et al. Tanshinone I attenuates the effects of a challenge with H<sub>2</sub>O<sub>2</sub>, on the functions of tricarboxylic acid cycle and respiratory chain in SH-SY5Y Cells [J]. *Mol Neurobiol*, 2017, **54**(10): 7858-7868.
- [51] Park JH, Park OK, Yan BC, et al. Neuroprotection via maintenance or increase of antioxidants and neurotrophic factors in ischemic gerbil hippocampus treated with tanshinone I [J]. *Chin Med J (Engl)*, 2014, **127**(19): 3396-3405.
- [52] Chen CP, Yokozawa T, Chung HY. Inhibitory effect of caffeic acid analogues isolated from *Salviae miltiorrhizae* Radix against 1, 1-diphenyl-2-picrylhydrazyl radical [J]. *Exp Toxicol Pathol*, 1999, **51**(1): 59-63.
- [53] Wu CF, Bohnert S, Thines E, et al. Cytotoxicity of *Salvia miltiorrhiza* against multidrug-resistant cancer cells [J]. *Am J Chin Med*, 2016, **44**(4): 871-894.
- [54] Chen XP, Guo JJ, Bao JL, et al. The anticancer properties of *Salvia miltiorrhiza* bunge (Danshen): a systematic review [J]. *Med Res Rev*, 2014, **34**(4): 768-794.
- [55] Kim EO, Kang SE, Im CR, et al. Tanshinone IIA induces TRAIL sensitization of human lung cancer cells through selective ER stress induction [J]. *Int J Oncol*, 2016, **48**(5): 2205-2212.
- [56] Yu J, Wang XX, Li YH, et al. Tanshinone IIA suppresses gastric cancer cell proliferation and migration by downregulation of FOXM1 [J]. *Oncol Rep*, 2017, **37**(3): 1394-1400.
- [57] Zhang HS, Zhang FJ, Li H, et al. Tanshinone IIA inhibits human esophageal cancer cell growth through miR-122-mediated PKM2 down-regulation [J]. *Arch Biochem Biophys*, 2016, **598**: 50-56.
- [58] Ding L, Wang S, Qu X, et al. Tanshinone IIA sensitizes oral squamous cell carcinoma to radiation due to an enhanced autophagy [J]. *Environ Toxicol Pharmacol*, 2016, **46**: 264-269.
- [59] Sung HJ, Choi SM, Yoon Y, et al. Tanshinone IIA, an ingredient of *Salvia miltiorrhiza* BUNGE, induces apoptosis in human leukemia cell lines through the activation of caspase-3 [J]. *Exp Mol Med*, 1999, **31**(4): 174-178.
- [60] Lee CY, Sher HF, Chen HW, et al. Anticancer effects of tanshinone I in human non-small cell lung cancer [J]. *Mol Cancer Ther*, 2008, **7**(11): 3527-3538.
- [61] Jing X, Xu Y, Cheng W, et al. Tanshinone I induces apoptosis and pro-survival autophagy in gastric cancers [J]. *Cancer Chemother Pharmacol*, 2016, **77**(6): 1171-1181.
- [62] Wang L, Wu JZ, Lu JW, et al. Regulation of the cell cycle and PI3K/Akt/mTOR signaling pathway by tanshinone I in human breast cancer cell lines [J]. *Mol Med Rep*, 2015, **11**(2): 931-939.
- [63] Shin EA, Sohn EJ, Won G, et al. Upregulation of microRNA-135a-3p and death receptor 5 plays a critical role in tanshinone I sensitized prostate cancer cells to TRAIL induced apoptosis [J]. *Oncotarget*, 2014, **5**(14): 5624-5636.
- [64] Chen WX, Pan YH, Wang SL, et al. Cryptotanshinone activates AMPK-TSC2 axis leading to inhibition of mTORC1 signaling in cancer cells [J]. *Bmc Cancer*, 2017, **17**(1): 34-44.
- [65] Chen XP, Li QH, He Y, et al. 15, 16-dihydrotanshinone I induces apoptosis and inhibits the proliferation, migration of human osteosarcoma cell line 143B *in vitro* [J]. *Anticancer Agents Med Chem*, 2017, **17**(9): 1234-1242.

- [66] Zhang Q, Wang SF, Yu YY, et al. Salvianolic acid A, as a novel ETA receptor antagonist, shows inhibitory effects on tumor *in vitro* [J]. *Int J Mol Sci*, 2016, **17**(8): E1244.
- [67] Li H, Shi L, Wei J, et al. Cellular uptake and anticancer activity of salvianolic acid B phospholipid complex loaded nanoparticles in head and neck cancer and precancer cells [J]. *Colloids Surf B Biointerfaces*, 2016, **147**: 65-72.
- [68] Wang X, Wang CY, Zhang LJ, et al. Salvianolic acid A shows selective cytotoxicity against multidrug-resistant MCF-7 breast cancer cells [J]. *Anticancer Drugs*, 2015, **26**(2): 210-223.
- [69] Wu CF, Karioti A, Rohr D, et al. Production of rosmarinic acid and salvianolic acid B from callus culture of *Salvia miltiorrhiza* with cytotoxicity towards acute lymphoblastic leukemia cells [J]. *Food Chem*, 2016, **201**: 292-297.
- [70] Wang ZS, Luo P, Dai SH, et al. Salvianolic acid B induces apoptosis in human glioma U87 cells through p38-mediated ROS generation [J]. *Cell Mol Neurobiol*, 2013, **33**(7): 921-928.
- [71] Chen Z, Xu H. Anti-inflammatory and immunomodulatory mechanism of tanshinone IIA for atherosclerosis [J]. *Evid Based Complement Alternat Med*, 2014, **2014**: 267976-267976.
- [72] Niu XL, Ichimori K, Yang X, et al. Tanshinone II-A inhibits low density lipoprotein oxidation *in vitro* [J]. *Free Radic Res*, 2000, **33**(3): 305-312.
- [73] Liu Z, Xu S, Huang X, et al. Cryptotanshinone, an orally bioactive herbal compound from Danshen, attenuates atherosclerosis in apolipoprotein E-deficient mice: role of LOX-1 [J]. *Br J Pharmacol*, 2015, **172**(23): 5661-5675.
- [74] Zhao W, Wu C, Chen X. Cryptotanshinone inhibits oxidized LDL-induced adhesion molecule expression ROS dependent NF- $\kappa$ B pathways [J]. *Cell Adh Migr*, 2016, **10**(3): 248-258.
- [75] Ran X, Zhao W, Li W, et al. Cryptotanshinone inhibits TNF- $\alpha$ -induced LOX-1 expression by suppressing reactive oxygen species (ROS) formation in endothelial cells [J]. *Korean J Physiol Pharmacol*, 2016, **20**(4): 347-355.
- [76] Zhao W, Li C, Gao H, et al. Dihydro-tanshinone I attenuates atherosclerosis in apoE-deficient mice: role of NOX4/NF- $\kappa$ B mediated lectin-like oxidized LDL receptor-1 (LOX-1) of the endothelium [J]. *Front Pharmacol*, 2016, **7**(117): 418.
- [77] Cho YH, Ku CR, Hong ZY, et al. Therapeutic effects of water soluble Danshen extracts on atherosclerosis [J]. *Evid Based Complement Alternat Med*, 2013, **2013**: 623639-623639.
- [78] Rang, Humphrey. *Rang and Dale's Pharmacology* [M]. ELSEVIER, 2007: 156-158.
- [79] Cao H, Zhang L, Sun ZB, et al. *Salvia miltiorrhiza* prevents deep vein thrombosis *via* antioxidative effects in endothelial cells [J]. *Mol Med Rep*, 2015, **11**(5): 3593-3600.
- [80] Lü M, Wang TY, Tian XX, et al. Interaction of anti-thrombotic and anti-inflammatory activities of commonly used traditional Chinese medicine for promoting blood circulation and removing blood stasis revealed by network pharmacology analysis [J]. *Acta Pharm Sin*, 2015, **50**(9): 1135-1141.
- [81] Huang ZS, Zeng CL, Zhu LJ, et al. Salvianolic acid A inhibits platelet activation and arterial thrombosis *via* inhibition of phosphoinositide 3-kinase [J]. *J Thromb Haemost*, 2010, **8**(6): 1383-1393.
- [82] Moon CY, Ku CR, Cho YH, et al. Protocatechuic aldehyde inhibits migration and proliferation of vascular smooth muscle cells and intravascular thrombosis [J]. *Biochem Biophys Res Commun*, 2012, **423**(1): 116-121.
- [83] Rang, Humphrey. *Rang and Dale's Pharmacology* [M]. ELSEVIER, 2007: 231-235.
- [84] Wang Y, Cao SH, Cui YJ, et al. *Salvia miltiorrhiza* Bge.f.alba ameliorates the progression of monocrotaline-induced pulmonary hypertension by protecting endothelial injury in rats [J]. *Tohoku J Exp Med*, 2015, **236**(2): 155-162.
- [85] Kang DG, Yun YG, Ryou JH, et al. Anti-hypertensive effect of water extract of danshen on renovascular hypertension through inhibition of the renin angiotensin system [J]. *Am J Chin Med*, 2002, **30**(01): 87-93.
- [86] Zhang J, An SJ, Fu JQ, et al. Mixed aqueous extract of *Salvia miltiorrhiza* reduces blood pressure through inhibition of vascular remodelling and oxidative stress in spontaneously hypertensive rats [J]. *Cell Physiol Biochem*, 2016, **40**(1-2): 347-360.
- [87] Wang J, Lu WJ, Wang W, et al. Promising therapeutic effects of sodium tanshinone IIA sulfonate towards pulmonary arterial hypertension in patients [J]. *J Thorac Dis*, 2013, **5**(2): 169-172.
- [88] Ling WC, Liu J, Lau CW, et al. Treatment with salvianolic acid B restores endothelial function in angiotensin II-induced hypertensive mice [J]. *Biochem Pharmacol*, 2017, **136**: 76-85.
- [89] Chen J, Deng J, Zhang YY, et al. Lipid-lowering effects of Danhong injection on hyperlipidemia rats [J]. *J Ethnopharmacology*, 2014, **154**(2): 437-442.
- [90] Zhang SJ, Cheng ZX, Lin YW, et al. Efficacy of composite salviae dropping pill on hyperlipemia patients with phlegm and blood stasis syndrome [J]. *Chin J Chin Mater Med*, 2007, **32**(5): 440-443.
- [91] Jia LQ, Zhang N, Xu Y, et al. Tanshinone IIA affects the HDL subfractions distribution not serum lipid levels: involving in intake and efflux of cholesterol [J]. *Arch Biochem Biophys*, 2016, **592**: 50-59.
- [92] Rahman N, Jeon M, Song HY, et al. Cryptotanshinone, a compound of *Salvia miltiorrhiza* inhibits pre-adipocytes differentiation by regulation of adipogenesis-related genes expression via STAT3 signaling [J]. *Phytomedicine*, 2016, **23**(1): 58-67.
- [93] Yang QA, Wang SW, Xie YH, et al. Effect of salvianolic acid b and paeonol on blood lipid metabolism and hemorrheology in myocardial ischemia rabbits induced by pituitrin [J]. *Int J Mol Sci*, 2010, **11**(10): 3696-3704.
- [94] Xu GL, Zhao WX, Zhou ZM, et al. Danshen extracts decrease blood C reactive protein and prevent ischemic stroke recurrence: a controlled pilot study [J]. *Phytother Res*, 2009, **23**(12): 1721-1725.
- [95] Lv HD, Wang L, Shen JC, et al. Salvianolic acid B attenuates apoptosis and inflammation *via* SIRT1 activation in experimental stroke rats [J]. *Brain Res Bull*, 2015, **115**: 30-36.
- [96] Tzen JT, Jinn TR, Chen YC, et al. Magnesium lithospermate B possesses inhibitory activity on Na<sup>+</sup>, K<sup>+</sup>-ATPase and neuroprotective effects against ischemic stroke [J]. *Acta Pharmacol Sin*, 2007, **28**(5): 609-615.
- [97] Li DC, Bao XQ, Sun H, et al. Research progress in the study of protective effect of tanshinone IIA on cerebral ischemic stroke [J].

*Acta Pharm Sin*, 2015, **50**(6): 635-639.

- [98] Yu XY, Lin SG, Chen X, et al. Transport of cryptotanshinone, a major active triterpenoid in *Salvia miltiorrhiza* Bunge widely used in the treatment of stroke and Alzheimer's disease, across the blood-brain barrier [J]. *Curr Drug Metab*, 2007, **8**(4): 365-377.
- [99] Liu BY, Du YH, Cong LX, et al. Danshen (*Salvia miltiorrhiza*) compounds improve the biochemical indices of the patients with coronary heart disease [J]. *Evid Based Complement Alternat Med*, 2016, **2016**: 9781715-9781715.
- [100] Huang JC, Tang XJ, Ye FX, et al. Clinical therapeutic effects of aspirin in combination with Fufang Danshen Diwan, a traditional Chinese medicine formula, on coronary heart disease: a systematic review and meta-analysis[J]. *Cell Physiol Biochem*, 2016, **39**(5): 1955-1963.
- [101] Wei W, Liu YW, Zhang Q, et al. Danshen-enhanced cardioprotective effect of cardioplegia on ischemia reperfusion injury in a human-induced pluripotent stem cell-derived cardiomyocytes model [J]. *Artif Organs*, 2017, **41**(5): 452-460.
- [102] Qin RA, Lin J, Li CY, et al. Study of the protective mechanisms of compound Danshen Tablet (Fufang Danshen Pian) against myocardial ischemia/reperfusion injury via the Akt-eNOS signaling pathway in rats [J]. *J Ethnopharmacol*, 2014, **156**: 190-198.
- [103] Li Q, Shen L, Wang Z, et al. Tanshinone IIA protects against myocardial ischemia reperfusion injury by activating the PI3K/Akt/mTOR signaling pathway [J]. *Biomed Pharmacother*, 2016, **84**: 106-114.
- [104] Wei Y, Xu M, Ren Y, et al. The cardioprotection of dihydrotanshinone I against myocardial ischemia-reperfusion injury via inhibition of arachidonic acid  $\omega$ -hydroxylase [J]. *Can J Physiol Pharmacol*, 2016, **94**(12): 1267-1275.
- [105] Yu JH, Wang LY, Akinyi M, et al. Danshensu protects isolated heart against ischemia reperfusion injury through activation of Akt/ERK1/2/Nrf2 signaling [J]. *Int J Clin Exp Med*, 2015, **8**(9): 14793-14804.
- [106] He ZF, Sun CZ, Xu Y, et al. Reduction of atrial fibrillation by Tanshinone IIA in chronic heart failure [J]. *Biomed Pharmacother*, 2016, **84**: 1760-1767.
- [107] Huang LB, Zheng M, Zhou YD, et al. Tanshinone IIA attenuates cardiac dysfunction in endotoxin-induced septic mice via inhibition of NADPH oxidase 2-related signaling pathway [J]. *Int Immunopharmacol*, 2015, **28**(1): 444-449.
- [108] Lee CM, Wong HN, Chui KY, et al. Miltirone, a central benzodiazepine receptor partial agonist from a Chinese medicinal herb *Salvia miltiorrhiza* [J]. *Neurosci Lett*, 1991, **127**(2): 237-241.
- [109] Fang XS, Hao JF, Zhou HY, et al. Pharmacological studies on the sedative-hypnotic effect of Semen Ziziphi spinosae (Suanzaoren) and Radix et Rhizoma *Salviae miltiorrhizae* (Danshen) extracts and the synergistic effect of their combinations [J]. *Phytomedicine*, 2010, **17**(1): 75-80.
- [110] Liu C, Shi W, Sun L, et al. Effects of radix *Salviae miltiorrhizae* on visceral pain discharges in the posterior nucleus of the thalamus in cats [J]. *Chin J Chin Mater Med*, 1990, **15**(2): 112-115.
- [111] Cao FL, Xu M, Wang Y, et al. Tanshinone IIA attenuates neuropathic pain via inhibiting glial activation and immune response [J]. *Pharmacol Biochem Behav*, 2015, **128**: 1-7.
- [112] Tang J, Zhu C, Li ZH, et al. Inhibition of the spinal astrocytic JNK/MCP-1 pathway activation correlates with the analgesic effects of tanshinone IIA sulfonate in neuropathic pain [J]. *J Neuroinflammation*, 2015, **12**(1):57.
- [113] Li J, Wen PY, Li WW, et al. Upregulation effects of Tanshinone IIA on the expressions of NeuN, Nissl body, and I $\kappa$ B and downregulation effects on the expressions of GFAP and NF- $\kappa$ B in the brain tissues of rat models of Alzheimer's disease [J]. *Neuroreport*, 2015, **26**(13): 758-766.
- [114] Wen P, Luo H, Zhou L, et al. Effects of tanshinone IIA on the expressions of caspase-3, Akt and NF- $\kappa$ B in the brains of rat models of Alzheimer's disease [J]. *Chin J Cell Mol Imm*, 2014, **30**(2): 155-159.
- [115] Jiang P, Li C, Xiang Z, et al. Tanshinone IIA reduces the risk of Alzheimer's disease by inhibiting iNOS, MMP-2 and NF- $\kappa$ B p65 transcription and translation in the temporal lobes of rat models of Alzheimer's disease [J]. *Mol Med Rep*, 2014, **10**(2): 689-694.
- [116] Mei Z, Yan P, Situ B, et al. Cryptotanshinone inhibits  $\beta$ -amyloid aggregation and protects damage from  $\beta$ -amyloid in SH-SY5Y cells [J]. *Neurochem Res*, 2012, **37**(3), 622-628.
- [117] Mei ZR, Zhang FY, Tao L, et al. Cryptotanshinone, a compound from *Salvia miltiorrhiza* modulates amyloid precursor protein metabolism and attenuates beta-amyloid deposition through upregulating alpha-secretase *in vivo* and *in vitro* [J]. *Neurosci Lett*, 2009, **452**(2): 90-95.
- [118] Wong K, Ho T, Lin HQ, et al. Cryptotanshinone, an acetylcholinesterase inhibitor from *Salvia miltiorrhiza*, ameliorates scopolamine-induced amnesia in Morris water maze task [J]. *Planta Med*, 2010, **76**(3): 228-234.
- [119] Wong KKK, Ngo JCK, Liu SJ, et al. Interaction study of two diterpenes, cryptotanshinone and dihydrotanshinone, to human acetylcholinesterase and butyrylcholinesterase by molecular docking and kinetic analysis [J]. *Chem Biol Interact*, 2010, **187**(1-3): 335-339.
- [120] Zhou YQ, Li WX, Xu L, et al. In *Salvia miltiorrhiza*, phenolic acids possess protective properties against amyloid beta-induced cytotoxicity and tanshinones act as acetylcholinesterase inhibitors [J]. *Environ Toxicol Pharmacol*, 2011, **31**(3): 443-452.
- [121] Cao YY, Wang L, Ge H, et al. Salvianolic acid A, a polyphenolic derivative from *Salvia miltiorrhiza* bunge, as a multifunctional agent for the treatment of Alzheimer's disease [J]. *Mol Divers*, 2013, **17**(3): 515-524.
- [122] Lee YW, Kim DH, Jeon SJ, et al. Neuroprotective effects of salvianolic acid B on an A $\beta$ 25-35 peptide-induced mouse model of Alzheimer's disease [J]. *Eur J Pharmacol*, 2013, **704**(1-3): 70-77.
- [123] Tang Y, Huang D, Zhang MH, et al. Salvianolic acid B inhibits a beta generation by modulating BACE1 activity in SH-SY5Y-APPsw cells [J]. *Nutrients*, 2016, **8**(6): E333.
- [124] Chong CM, Zhou ZY, Razmovski-Naumovski V, et al. Danshensu protects against 6-hydroxydopamine-induced damage of PC12 cells *in vitro* and dopaminergic neurons in zebrafish [J].

- Neurosci Lett*, 2013, **543**: 121-125.
- [125] Zhou J, Qu XD, Li ZY, et al. Salvianolic acid B attenuates toxin-induced neuronal damage via Nrf2-dependent glial cell-mediated protective activity in Parkinson's disease models [J]. *PLoS One*, 2014, **9**(7): e101668.
- [126] Ji K, Zhao Y, Yu T, et al. Inhibition effects of tanshinone on the aggregation of  $\alpha$ -synuclein [J]. *Food Funct*, 2015, **7**(1): 409-416.
- [127] Wang S, Jing H, Yang H, et al. Tanshinone I selectively suppresses pro-inflammatory genes expression in activated microglia and prevents nigrostriatal dopaminergic neurodegeneration in a mouse model of Parkinson's disease [J]. *J Ethnopharmacol*, 2015, **164**: 247-255.
- [128] Xu J, Wei X, Ren M, et al. Neuroprotective effects of Tanshinone I against 6-OHDA-induced oxidative stress in cellular and mouse model of Parkinson's disease through upregulating Nrf2 [J]. *Neurochem Res*, 2015, **41**(4): 779-786.
- [129] Ren B, Zhang YX, Zhou HX, et al. Tanshinone IIA prevents the loss of nigrostriatal dopaminergic neurons by inhibiting NADPH oxidase and iNOS in the MPTP model of Parkinson's disease [J]. *J Neurol Sci*, 2015, **348**(1-2): 142-152.
- [130] Li L. Protective effects of schisanhenol, salvianolic acid A and SY-L on oxidative stress induced injuries of cerebral cells and their mechanisms [J]. *Prog Physiol Sci*, 1998, **29**(1): 35-38.
- [131] Tian LL, Wang XJ, Sun YN, et al. Salvianolic acid B, an antioxidant from *Salvia miltiorrhiza*, prevents 6-hydroxydopamine induced apoptosis in SH-SY5Y cells [J]. *Int J Biochem Cell Biol*, 2008, **40**(3): 409-422.
- [132] Zhang Y, Zhang YY, Xie Y, et al. Multitargeted inhibition of hepatic fibrosis in chronic iron-overloaded mice by *Salvia miltiorrhiza* [J]. *J Ethnopharmacol*, 2013, **148**(2): 671-681.
- [133] Nan JX, Park EJ, Kang HC, et al. Anti-fibrotic effects of a hot-water extract from *Salvia miltiorrhiza* roots on liver fibrosis induced by biliary obstruction in rats [J]. *J Pharm Pharmacol*, 2001, **53**(2): 197-204.
- [134] Sferri R, Vetuschi A, Catitti V, et al. Boswellia serrata and *Salvia miltiorrhiza* extracts reduce DMN-induced hepatic fibrosis in mice by TGF- $\beta$ 1 downregulation [J]. *Eur Rev Med Pharmacol Sci*, 2012, **16**(11): 1484-1498.
- [135] Wang WW, Guan CW, Sun XZ, et al. Tanshinone IIA protects against acetaminophen-induced hepatotoxicity via activating the Nrf2 pathway [J]. *Phytomedicine*, 2016, **23**(6): 589-596.
- [136] Jin Q, Jiang S, Wu YL, et al. Hepatoprotective effect of cryptotanshinone from *Salvia miltiorrhiza* in d-galactosamine/lipopolysaccharide-induced fulminant hepatic failure [J]. *Phytomedicine*, 2014, **21**(2): 141-147.
- [137] Ge M, Liu H, Zhang Y, et al. The anti-hepatic fibrosis effects of dihydrotanshinone I are mediated by disrupting the YAP and TEAD2 complex and stimulating autophagy [J]. *Br J Pharmacol*, 2017, **174**(10): 1147-1160.
- [138] Lv T, Yao XX. Comparison of protocatechuic aldehyde in Radix *Salvia miltiorrhiza* and corresponding pharmacological sera from normal and fibrotic rats by high performance liquid chromatography [J]. *World J Gastroenterol*, 2006, **12**(14): 2195-2200.
- [139] Ding CC, Zhao Y, Shi X, et al. New insights into salvianolic acid A action: regulation of the TXNIP/NLRP3 and TXNIP/ChREBP pathways ameliorates HFD-induced NAFLD in rats [J]. *Sci Rep*, 2016, **6**: 28734.
- [140] Lv Z, Xu L. Salvianolic acid B inhibits ERK and p38 MAPK signaling in TGF- $\beta$ 1-stimulated human hepatic stellate cell line (LX-2) via distinct pathways [J]. *Evid Based Complement Alternat Med*, 2012, **2012**(4): 960128.
- [141] Paik YH, Yoon YJ, Lee HC, et al. Antifibrotic effects of magnesium lithospermate B on hepatic stellate cells and thioacetamide-induced cirrhotic rats [J]. *Exp Mol Med*, 2011, **43**(6): 341-349.
- [142] Chan KWK, Ho WS. Anti-oxidative and hepatoprotective effects of lithospermic acid against carbon tetrachloride-induced liver oxidative damage *in vitro* and *in vivo* [J]. *Oncol Rep*, 2015, **34**(2): 673-680.
- [143] Dai LJ, Hou J, Cai HR. Experimental study on treatment of pulmonary fibrosis by Chinese drugs and integrative Chinese and Western medicine [J]. *Chin J Integr Trad West Med*, 2004, **24**(2): 130-132.
- [144] Hou J, Dai L, Huang M, et al. The therapeutic effect of ligustrazin and *Salvia miltiorrhiza* on the gene expression of  $\alpha$ 1 (I) and  $\alpha$ 1 (III) procollagen in rat pulmonary fibrosis [J]. *Chin J Tuberc Respir Dis*, 1999, **22**(1): 43-45.
- [145] He HY, Tang HY, Gao LL, et al. Tanshinone IIA attenuates bleomycin-induced pulmonary fibrosis in rats [J]. *Mol Med Rep*, 2015, **11**(6): 4190-4196.
- [146] Pan YM, Fu HY, Kong Q, et al. Prevention of pulmonary fibrosis with salvianolic acid A by inducing fibroblast cell cycle arrest and promoting apoptosis [J]. *J Ethnopharmacol*, 2014, **155**(3): 1589-1596.
- [147] Zhang M, Cao SR, Zhang R, et al. The inhibitory effect of salvianolic acid B on TGF- $\beta$ 1-induced proliferation and differentiation in lung fibroblasts [J]. *Exp Lung Res*, 2014, **40**(4): 172-185.
- [148] Liu J, Hua G, Wang H, et al. Experimental study of the effect of IH764-3 on pulmonary fibrosis [J]. *Chin Med Sci J*, 1993, **8**(1): 9-14.
- [149] Yao DQ, Tian YP, Gao HB, et al. Study the effects of *Salvia miltiorrhiza* monomer IH764-3 on the expression of matrix metalloproteinase in lungs of rats exposed to Paraquat (PQ) [J]. *Chin J Ind Hyg Occup Dis*, 2012, **30**(5): 321-325.
- [150] Lu XM, Jin YN, Ma L, et al. Danshen (Radix *Salviae miltiorrhizae*) reverses renal injury induced by myocardial infarction [J]. *J Tradit Chin Med*, 2015, **35**(3): 306-311.
- [151] He L, Zhang QQ, Lü XY, et al. Effects of water extract of *Salvia miltiorrhiza* against renal injury on rats exposed to cadmium [J]. *Natl Med J China*, 2017, **97**(1): 57-61.
- [152] Chen G, Fu YR, Wu XH. Protective effect of *Salvia miltiorrhiza* extract against renal ischemia-reperfusion-induced injury in rats [J]. *Molecules*, 2012, **17**(2): 1191-1202.
- [153] Jiang CM, Shao QY, Jin B, et al. Tanshinone IIA attenuates renal fibrosis after acute kidney injury in a mouse model through inhibition of fibrocytes recruitment [J]. *Biomed Res Int*, 2015, **2015**: 867140-867140.
- [154] Jiang C, Zhu W, Yan X, et al. Rescue therapy with Tanshinone IIA hinders transition of acute kidney injury to chronic kidney

- disease via targeting GSK3 $\beta$  [J]. *Sci Rep*, 2016, **6**: 36698.
- [155] Xu YM, Ding GH, Huang J, et al. Tanshinone IIA pretreatment attenuates ischemia/reperfusion-induced renal injury [J]. *Exp Ther Med*, 2016, **12**(4): 2741-2746.
- [156] Kang DG, Oh H, Sohn EJ, et al. Lithospermic acid B isolated from *Salvia miltiorrhiza* ameliorates ischemia/reperfusion-induced renal injury in rats [J]. *Life Sci*, 2004, **75**(15): 1801-1816.
- [157] Park CH, Shin SH, Lee EK, et al. Magnesium lithospermate B from *Salvia miltiorrhiza* Bunge ameliorates aging-induced renal inflammation and senescence via NADPH oxidase-mediated reactive oxygen generation [J]. *Phytother Res*, 2017, **31**(5): 721-728.
- [158] Guan SJ, Ma JJ, Zhang Y, et al. Danshen (*Salvia miltiorrhiza*) injection suppresses kidney injury induced by iron overload in mice [J]. *PLoS One*, 2013, **8**(9): e74318.
- [159] Li L, Zhang YY, Ma JJ, et al. *Salvia miltiorrhiza* injection ameliorates renal damage induced by lead exposure in mice [J]. *Sci World J*, 2014, **2014**: 572697-572697.
- [160] Huang MQ, Xie YL, Chen LD, et al. Antidiabetic effect of the total polyphenolic acids fraction from *Salvia miltiorrhiza* Bunge in diabetic rats [J]. *Phytother Res*, 2012, **26**(6): 944-948.
- [161] Yang XY, Sun L, Xu P, et al. Effects of salvianolic acid A on plantar microcirculation and peripheral nerve function in diabetic rats [J]. *Eur J Pharmacol*, 2011, **665**(1-3): 40-46.
- [162] Raoufi S, Baluchnejadmojarad T, Roghani M, et al. Anti-diabetic potential of salvianolic acid B in multiple low-dose streptozotocin-induced diabetes [J]. *Pharm Biol*, 2015, **53**(12): 1803-1809.
- [163] Huang MQ, Zhou CJ, Zhang YP, et al. Salvianolic acid B ameliorates hyperglycemia and dyslipidemia in db/db mice through the AMPK pathway [J]. *Cell Physiol Biochem*, 2016, **40**(5): 933-943.
- [164] Ren YN, Tao SJ, Zheng SG, et al. Salvianolic acid B improves vascular endothelial function in diabetic rats with blood glucose fluctuations via suppression of endothelial cell apoptosis [J]. *Eur J Pharmacol*, 2016, **791**: 308-315.
- [165] Jin CJ, Yu SH, Wang XM, et al. The effect of lithospermic acid, an antioxidant, on development of diabetic retinopathy in spontaneously obese diabetic rats [J]. *PLoS One*, 2014, **9**(6): e98232.
- [166] Hu J, Li YL, Li ZL, et al. Chronic supplementation of paeonol combined with Danshensu for the improvement of vascular reactivity in the cerebral basilar artery of diabetic rats [J]. *Int J Mol Sci*, 2012, **13**(11): 14565-14578.
- [167] Wang T, Fu FH, Han B, et al. Danshensu ameliorates the cognitive decline in streptozotocin-induced diabetic mice by attenuating advanced glycation end product-mediated neuroinflammation [J]. *J Neuroimmunol*, 2012, **245**(1-2): 79-86.
- [168] Chen P, Chen J, Zheng Q, et al. Pioglitazone, extract of compound Danshen dripping pill, and quercetin ameliorate diabetic nephropathy in diabetic rats [J]. *J Endocrinol Invest*, 2013, **36**(6): 422-427.
- [169] Zhang M, Li X, Jiu G, et al. Effect of danshen injection on expression of platelet membrane glycoproteins in patients with type II diabetes mellitus [J]. *Chin Med Mat*, 2003, **26**(10): 738-740.
- [170] Xu LH, Shen PQ, Bi YL, et al. Danshen injection ameliorates STZ-induced diabetic nephropathy in association with suppression of oxidative stress, pro-inflammatory factors and fibrosis [J]. *Int Immunopharmacol*, 2016, **38**: 385-394.
- [171] Yin DK, Yin JJ, Yang Y, et al. Renoprotection of Danshen injection on streptozotocin-induced diabetic rats, associated with tubular function and structure [J]. *J Ethnopharmacol*, 2014, **151**(1): 667-674.
- [172] Zhang JL, Cui M, He Y, et al. Chemical fingerprint and metabolic fingerprint analysis of Danshen injection by HPLC-UV and HPLC-MS methods [J]. *J Pharm Biomed Anal*, 2005, **36**(5): 1029-1035.
- [173] Zhao D, Han DE, Li N, et al. Simultaneous determination of six phenolic constituents of Danshen injection in rat plasma by LC-ESI-MS and its application to a pharmacokinetic study [J]. *Eur J Mass Spectrom (Chichester)*, 2011, **17**(4): 395-403.
- [174] Abdelazem IS, Chen HS, Bates RB, et al. Isolation of two highly potent and non-toxic inhibitors of human immunodeficiency virus type 1 (HIV-1) integrase from *Salvia miltiorrhiza* [J]. *Antivir Res*, 2002, **55**(1): 91-106.
- [175] Zhang HS, Chen XY, Wu TC, et al. Tanshinone II A inhibits tat-induced HIV-1 transactivation through redox-regulated AMPK/Nampt pathway [J]. *J Cell Physiol*, 2014, **229**(9): 1193-1201.
- [176] Wang GZ, Ru X, Ding LH, et al. Short-term effect of *Salvia miltiorrhiza* in treating rat acetic acid chronic gastric ulcer and long-term effect in preventing recurrence [J]. *World J Gastroenterol*, 1998, **4**(2): 169-170.
- [177] Murakami S, Kijima H, Isohe Y, et al. Effect of salvianolic acid A, a depside from roots of *Salvia miltiorrhiza*, on gastric H<sup>+</sup>, K<sup>(+)</sup>-ATPase [J]. *Planta Med*, 1990, **56**(04): 360-363.
- [178] Ryu SY, Oak MH, Kim KM. Inhibition of mast cell degranulation by tanshinones from the roots of *Salvia miltiorrhiza* [J]. *Planta Med*, 1999, **65**(7): 654-655.
- [179] Trinh HT, Chae SJ, Joh EH, et al. Tanshinones isolated from the rhizome of *Salvia miltiorrhiza* inhibit passive cutaneous anaphylaxis reaction in mice [J]. *J Ethnopharmacol*, 2010, **132**(1): 344-348.
- [180] Yang JH, Son KH, Son JK, et al. Anti-allergic activity of an ethanol extract from *Salvia miltiorrhiza* [J]. *Arch Pharm Res*, 2008, **31**(12): 1597-1603.
- [181] Zhang CS, Yu JJ, Parker S, et al. Oral Chinese herbal medicine combined with pharmacotherapy for psoriasis vulgaris: a systematic review [J]. *Int J Dermatol*, 2014, **53**(11): 1305-1318.
- [182] Lee DS, Lee SH, Noh JG, et al. Antibacterial activities of cryptotanshinone and dihydrotanshinone I from a medicinal herb, *Salvia miltiorrhiza* Bunge [J]. *Biosci Biotechnol Biochem*, 1999, **63**(12): 2236-2239.
- [183] Liu QQ, Han J, Zuo GY, et al. Potentiation activity of multiple antibacterial agents by salvianolate from the Chinese medicine Danshen against methicillin-resistant staphylococcus aureus (MRSA) [J]. *J Pharmacol Sci*, 2016, **131**(1): 13-17.
- [184] Liang BF, Su JW. Involvement of renin-angiotensin system inhibition, the potential risk of Danshen in the treatment of pregnancy-induced hypertension [J]. *Phytother Res*, 2015, **29**(9): 1305-1318.

- 1421-1422.
- [185] Liu Y, Huang YH, Zhao CY, et al. *Salvia miltiorrhiza* injection on pulmonary heart disease: a systematic review and meta-analysis [J]. *Phytother Res*, 2014, **42**(6): 1315-1331.
- [186] Chang YP, Zhang W, Xie YM, et al. Postmarketing safety evaluation: deposite salt injection made from Danshen (*Radix Salviae miltiorrhizae*) [J]. *J Tradit Chin Med*, 2014, **34**(6): 749-753.
- [187] Wang C, Zhao R, Li B, et al. An *in vivo* and *in vitro* study: high-dosage Danshen injection induces peripheral vascular endothelial cells injury [J]. *Hum Exp Toxicol*, 2016, **35**(4): 404-417.
- [188] Wang N, Luo HW, Niwa M, et al. A new platelet aggregation inhibitor from *Salvia miltiorrhiza* [J]. *Planta Med*, 1989, **55**(4): 390-391.
- [189] Lin HC, Ding HY, Chang WL. Two new fatty diterpenoids from *Salvia miltiorrhiza* [J]. *J Nat Prod*, 2001, **64**(5): 648-650.
- [190] Ikeshiro Y, Mase I, Tomita Y. Abietane type diterpenoids from *Salvia miltiorrhiza* [J]. *Phytochemistry*, 1989, **28**(11): 3139-3141.
- [191] Onitsuka M, Fujiu M, Shinma N, et al. New platelet aggregation inhibitors from Tan-Shen; radix of *Salvia miltiorrhiza* Bunge [J]. *Chem Pharm Bull (Tokyo)*, 1983, **31**(5): 1670-1675.
- [192] Don MJ, Shen CC, Syu WJ, et al. Cytotoxic and aromatic constituents from *Salvia miltiorrhiza* [J]. *Phytochemistry*, 2006, **67**(5): 497-503.
- [193] Nagy G, Gunther G, Mathe I, et al. Diterpenoids from *Salvia glutinosa*, *S-austriaca*, *S-tomentosa* and *S-verticillata* roots [J]. *Phytochemistry*, 1999, **52**(6): 1105-1109.
- [194] Li YG, Song L, Liu M, et al. Advancement in analysis of *Salviae miltiorrhizae* Radix et Rhizoma (Danshen) [J]. *J Chromatogr A*, 2009, **1216**(11): 1941-1953.
- [195] Li HB, Chen F. Preparative isolation and purification of salidroside from the Chinese medicinal plant *Rhodiola sachalinensis* by high-speed counter-current chromatography [J]. *J Chromatogr A*, 2001, **932**(1-2): 91-95.
- [196] Yang M, Liu A, Guan S, et al. Characterization of tanshinones in the roots of *Salvia miltiorrhiza* (Dan-shen) by high-performance liquid chromatography with electrospray ionization tandem mass spectrometry [J]. *Rapid Commun Mass Spectrom*, 2006, **20**(8): 1266-1280.
- [197] Ma ZJ, Zhang M, Song ZH. Characterization of tanshinones with quinone reductase induction activity from Radix *Salvia miltiorrhiza* by liquid chromatography/tandem mass spectrometry [J]. *Rapid Commun Mass Spectrom*, 2009, **23**(18): 2857-2866.
- [198] Lee AR, Wu WL, Chang WL, et al. Isolation and bioactivity of new tanshinones [J]. *J Nat Prod*, 1987, **50**(2): 157-160.
- [199] Han YM, Oh H, Na MK, et al. PTP1B inhibitory effect of abietane diterpenes isolated from *Salvia miltiorrhiza* [J]. *Biol Pharm Bull*, 2005, **28**(9): 1795-1797.
- [200] Luo HW, Wu BJ, Wu MY, et al. Pigments from *Salvia miltiorrhiza* [J]. *Phytochemistry*, 1985, **24**(4): 815-817.
- [201] Danheiser RL, Casebier DS, Loebach JL. Total synthesis of danshen diterpenoid quinones [J]. *Tetrahedron Lett*, 1992, **33**(9): 1149-1152.
- [202] Luo HW, Wu BJ, Wu MY, et al. Isolation and structure of danshenxinkun D [J]. *Acta Pharm Sin*, 1985, **20**(7): 542-544.
- [203] Yu XY, Lin SG, Zhou ZW, et al. Tanshinone IIB, a primary active constituent from *Salvia miltiorrhiza*, exhibits neuroprotective activity in experimentally stroked rats [J]. *Neurosci Lett*, 2007, **417**(3): 261-265.
- [204] Dat NT, Jin X, Lee JH, et al. Abietane diterpenes from *Salvia miltiorrhiza* inhibit the activation of hypoxia-inducible factor-1 [J]. *J Nat Prod*, 2007, **70**(7): 1093-1097.
- [205] Wang XH, Morris-Natschke SL, Lee KH. New developments in the chemistry and biology of the bioactive constituents of Tanshen [J]. *Med Res Rev*, 2007, **27**(1): 133-148.
- [206] Xuezhao L, Houwei L, Masatake N. Trijuganone A and B: two new phenanthrenequinones from roots of *Salvia trijuga* [J]. *Planta Med*, 1990, **56**(1): 87-88.
- [207] Lin HC, Chang WL. Diterpenoids from *Salvia miltiorrhiza* [J]. *Phytochemistry*, 2000, **53**(8): 951-953.
- [208] Ryu SY, No Z, Kim SH, et al. Two novel abietane diterpenes from *Salvia miltiorrhiza* [J]. *Planta Med*, 1997, **63**(1): 44-46.
- [209] Wang XH, Bastow KF, Sun CM, et al. Antitumor agents. 239. isolation, structure elucidation, total synthesis, and anti-breast cancer activity of neo-tanshinolactone from *Salvia miltiorrhiza* [J]. *J Med Chem*, 2004, **47**(23): 5816-5819.
- [210] Yang BJ, Qian MK, Qin GW, et al. Studies on the active principles of Dan-Shen. V. isolation and structures of prezewaquinone A and prezewaquinone B (author's transl) [J]. *Acta Pharm Sin*, 1981, **16**(11): 837-841.
- [211] Don MJ, Shen CC, Lin YL, et al. Nitrogen-containing compounds from *Salvia miltiorrhiza* [J]. *J Nat Prod*, 2005, **68**(7): 1066-1070.
- [212] Li XX, Xu X, Wang JN, et al. A system-level investigation into the mechanisms of Chinese Traditional Medicine: compound Danshen Formula for cardiovascular disease treatment [J]. *PLoS One*, 2012, **7**(9): e43918.
- [213] Asari F, Kusumi T, Zheng GZ, et al. Cryptoceralide and epicryptoceralide, novel spiro lactone diterpenoids from *Salvia miltiorrhiza* [J]. *Chem Lett*, 1990(10): 1885-1888.
- [214] Lin HC, Chang WL. Phytochemical and pharmacological study on *Salvia miltiorrhiza* (VI)-cytotoxic activity of tanshinones [J]. *Chin Pharm J*, 1995, **47**(1): 77-80.
- [215] Feng B, Li S. Studies on the chemical components of Dan-shen (*Salvia miltiorrhiza* Bunge) [J]. *Acta Pharm Sin*, 1980, **15**(8): 489-494.
- [216] Lin HC, Chang WL. Phytochemical and pharmacological study on *Salvia miltiorrhiza* (I)-isolation of new tanshinones [J]. *Chin Pharm J*, 1991, **43**(1): 11-17.
- [217] Kusumi T, Ooi T, Hayashi T, et al. A diterpenoid phenalenone from *Salvia miltiorrhiza* [J]. *Phytochemistry*, 1985, **24**(9): 2118-2120.
- [218] Liu J, Dai Z, Wang GL, et al. Progress in bioactive constituents and isolation and analysis methods of *Salvia Miltiorrhizae* Radix et Rhizoma [J]. *Chin J Exp Tradit Med Form*, 2012, **18**(11): 288-295.
- [219] Sun AL, Zhang YQ, Li AF, et al. Extraction and preparative purification of tanshinones from *Salvia miltiorrhiza* Bunge by high-speed counter-current chromatography [J]. *J Chromatogr*

- B Analyt Technol Biomed Life Sci*, 2011, **879**(21): 1899-1904.
- [220] Kuo YH, Wu CH. Synthesis of 5-(3-Hydroxypropyl)-7-methoxy-2-(3'-methoxy-4'-hydroxyphenyl)-3-benzo[b]furancarbaldehyde, a novel adenosine A1 receptor ligand from the root of *Salvia miltiorrhiza* [J]. *J Nat Prod*, 1996, **59**(6): 625-628.
- [221] Ikeshiro Y, Hashimoto I, Iwamoto Y, et al. Diterpenoids from *Salvia miltiorrhiza* [J]. *Phytochemistry*, 1991, **30**(8): 2791-2792.
- [222] Zhang HJ, Li LN. Salvianolic acid I: a new depside from *Salvia cavaleriei* [J]. *Planta Med*, 1994, **60**(1): 70-72.
- [223] Yang C, Zhang B. Extraction and isolation of water-soluble active constituent, Dan Shen Su, from *Salvia miltiorrhiza* and preparation of injections [J]. *Chin Pharm J*, 1981, **16**(11): 646-647.
- [224] Ai CB, Deng QH, Song WZ, et al. Salvianolic acid J, a depside from *Salvia flava* [J]. *Phytochemistry*, 1994, **37**(3): 907-908.
- [225] Kang HS, Chung HY, Jung JH, et al. Antioxidant effect of *Salvia miltiorrhiza* [J]. *Arch Pharm Res*, 1997, **20**(5): 496.
- [226] Chun Bo AI, Li LN. Synthesis of tetramethyl salvianolic acid F and (±)-trimethyl przewalskinic acid A [J]. *Chin Chem Lett*, 1996, **7**(5): 427-430.
- [227] Kohda H, Takeda O, Tanaka S, et al. Isolation of inhibitors of adenylate cyclase from dan-shen, the root of *Salvia miltiorrhiza* [J]. *Chem Pharm Bull (Tokyo)*, 1989, **37**(5): 1287-1290.
- [228] Yokozawa T, Chung HY, Oura H, et al. Isolation of a renal function-facilitating constituent from the Oriental drug, *Salviae miltiorrhizae* radix [J]. *Nihon Jinzo Gakkai Shi*, 1989, **31**(10): 1091-1098.
- [229] Hase K, Kasimu R, Basnet P, et al. Preventive effect of lithospermate B from *Salvia miltiorrhiza* on experimental hepatitis induced by carbon tetrachloride or D-galactosamine/lipopoly-saccharide [J]. *Planta Med*, 1997, **63**(1): 22.
- [230] Tanaka T, Morimoto S, Nonaka GI, et al. Magnesium and ammonium-potassium lithospermates B, the active principles having a uremia-preventive effect from *Salvia miltiorrhiza* [J]. *Chem Pharm Bull (Tokyo)*, 2008, **37**(2): 340-344.
- [231] Zhou CX, Luo HW, Niwa M. Studies on isolation and identification of water-soluble constituents of *Salvia miltiorrhiza* [J]. *J China Pharm Univ*, 1999, **30**(6): 411-416.
- [232] Li CBALN. Salvianolic acid G, a caffeic acid dimer with a novel tetracyclic skeleton [J]. *Chin Chem Lett*, 1991(1): 17-18.
- [233] Yang Z, Hon PM, Chui KY, et al. ChemInform abstract: naturally occurring benzofuran: isolation, structure elucidation, and total synthesis of 5-(3-Hydroxypropyl)-7-methoxy-2-(3'-methoxy-4'-hydroxyphenyl)-3-benzo(b)furancarbaldehyde, a novel adenosine A1 receptor ligand isolated from *Salvia miltiorrhiza* bunge (danshen) [J]. *Cheminform*, 1991, **32**(18): 2061-2064.
- [234] Chang JY, Chang CY, Kuo CC, et al. Salvinal, a novel microtubule inhibitor isolated from *Salvia miltiorrhizae* Bunge (Danshen), with antimetabolic activity in multidrug-sensitive and -resistant human tumor cells [J]. *Mol Pharmacol*, 2004, **65**(1): 77-84.
- [235] Lu Y, Foo LY. Salvianolic acid L, a potent phenolic antioxidant from *Salvia officinalis* [J]. *Cheminform*, 2001, **42**(46): 8223-8225.
- [236] Kohda H, Takeda O, Tanaka S, et al. Isolation of inhibitors of adenylate cyclase from Danshen, the root of *Salvia miltiorrhiza* [J]. *Chem Pharm Bull (Tokyo)*, 1989, **37**(5): 1287-1290.
- [237] Kasimu R, Tanaka K, Tezuka Y, et al. Comparative study of seventeen *Salvia* plants: aldose reductase inhibitory activity of water and MeOH extracts and liquid chromatography-mass spectrometry (LC-MS) analysis of water extracts [J]. *Chem Pharm Bull (Tokyo)*, 1998, **46**(3): 500-504.
- [238] Li W, Zhou SP, Jin YP, et al. Salvianolic acids T and U: a pair of atropisomeric trimeric caffeic acids derivatives from root of *Salvia miltiorrhiza* [J]. *Fitoterapia*, 2014, **98**: 248-253.
- [239] Zhu Z, Zhang H, Zhao L, et al. Rapid separation and identification of phenolic and diterpenoid constituents from radix *Salvia miltiorrhizae* by high-performance liquid chromatography diode-array detection, electrospray ionization time-of-flight mass spectrometry and electrospray ionization [J]. *Rapid Commun Mass Spectrom*, 2007, **21**(12): 1855-1865.
- [240] Li LN, Tan R, Chen WM. Salvianolic acid A, a new depside from roots of *Salvia miltiorrhiza* [J]. *Planta Med*, 1984, **50**(3): 227-228.
- [241] Qian L, Liang ZS, Wang JR, et al. Essential oil composition of *Salvia miltiorrhiza* flower [J]. *Food Chem*, 2009, **113**(2): 592-594.
- [242] Wang D, Girard TJ, Kasten TP, et al. Inhibitory activity of unsaturated fatty acids and anacardic acids toward soluble tissue factor-factor VIIa complex [J]. *J Nat Prod*, 1998, **61**(11): 1352-1355.
- [243] Ma C, Wang W, Chen YY, et al. Neuroprotective and antioxidant activity of compounds from the aerial parts of *dioscorea opposita* [J]. *J Nat Prod*, 2005, **68**(8): 1259-1261.
- [244] Rodríguez JA, Theoduloz C, Yáñez T, et al. Gastroprotective and ulcer healing effect of ferruginol in mice and rats: assessment of its mechanism of action using *in vitro* models [J]. *Life Sci*, 2006, **78**(21): 2503-2509.
- [245] Yang ZJ, Zhang L, Yang RW, et al. Principal component and cluster analysis of trace elements in Chinese herb *Salvia miltiorrhiza* and its relative species [J]. *Spectrosc Spect Anal*, 2008, **28**(10): 2441-2445.
- [246] Li H, Song F, Zhong Z, et al. Characterization of saccharides and phenolic acids in the Chinese herb Tanshen by ESI-FT-ICR-MS and HPLC [J]. *J Mass Spectrom*, 2008, **43**(11): 1545-1552.
- [247] Jr AJD, Michael W, Cecilia G. *Salvia columbariae* contains tanshinones [J]. *Evid Based Complement Alternat Med: eCAM*, 2005, **2**(1): 107-110.

**Cite this article as:** MEI Xiao-Dan, CAO Yan-Feng, CHE Yan-Yun, LI Jing, SHANG Zhan-Peng, ZHAO Wen-Jing, QIAO Yan-Jiang, ZHANG Jia-Yu. Danshen: a phytochemical and pharmacological overview [J]. *Chin J Nat Med*, 2018, **17**(1): 59-80.